

REGULAR SEIFERT SURFACES AND VASSILIEV KNOT INVARIANTS

EFSTRATIA KALFAGIANNI AND XIAO-SONG LIN

CONTENTS

Introduction

§1. Vassiliev knot invariants and Gusarov's n -triviality

§2. Regular Seifert surfaces of a knot

- 2a. Generalities
- 2b. Good position of a band in a regular spine
- 2c. Lower central series and curves on surfaces
- 2d. An example illustrating the good position of a band

§3. Commutator half bases and Vassiliev invariants

- 3a. Definitions and the main result
- 3b. Nice arcs and simple commutators
- 3c. Bad sets and good arcs
- 3d. Conflict sets and products of good arcs
- 3e. The reduction to nice arcs

§4. Vassiliev invariants as obstructions to n -sliceness

§5. More types of n -trivial knots

- 5a. Definitions
- 5b. Mixed type knots

§6. Discussion and questions

The first author was supported by NSF grant DMS-9626140, and the second author by NSF grants DMS-9400806 and DMS-9796130.

1991 Mathematics Subject Classification: 57M25.

This version: November 1999.

References

INTRODUCTION

It is well known ([Bi], [BN1]) that the Vassiliev knot invariants (or knot invariants of finite type) give a coherent and systematic way to view all the polynomial knot invariants (quantum invariants). Although both classes of the invariants have been studied extensively in the recent years, the following fundamental question is open: What geometric knot properties are Vassiliev invariants detecting? In particular to what extent does the vanishing of Vassiliev invariants approximate the unknottedness of a knot?

In the present paper we address the first question mentioned above. We show that the invariants are obstructions to a knot's bounding a *regular* Seifert surface whose complement looks, modulo the lower central series of its fundamental group, like the complement of a null-isotopy. See Theorems 3.2 and 5.4.

Milnor's $\bar{\mu}$ -invariants ([Mi1], [Mi2]) are integer link concordance invariants defined in terms of algebraic data extracted from the link group. By [BN2], [L2] and [HM], these invariants are the only link concordance invariants that are of finite type. Thus they can be thought of as the link homotopy (or link concordance) counterpart of Vassiliev's theory. As shown in [Co] and [O], Milnor's invariants detect exactly when the complements of links are cobordant as manifolds by a cobordism which looks, modulo the lower central series of its fundamental group, like the complement of a link concordance. Our main results here are parallel to results of [Co] and [O] and indicate that the two classes of invariants can be put under a unifying geometric framework.

The geometric notion that captures the vanishing of Milnor's invariants is the notion of *n-sliceness* or *n-null-concordance* ([Co],[O]). Before we state the main results of this paper, let us introduce some notation and terminology. We will say that a Seifert surface S of a knot K is *regular* if it has a spine Σ whose embedding in S^3 , induced by the embedding $S \subset S^3$, is isotopic to the standard embedding of a bouquet of circles. Such a spine will be called a *regular spine* of S . In particular, $\pi = \pi_1(S^3 \setminus S)$ is a free group. For each circle in a regular spine, we may push it off the surface S slightly in the normal directions. If these push-offs are all disjoint (this can be achieved by either pushing circles off to

different sides of S or in different distances away from S), we get a link $L^{\bar{\epsilon}}$. For “appropriate” $L^{\bar{\epsilon}}$ (see before the statement of Theorem 4.3 for details) we obtain that the Vassiliev invariants of the boundary knot $K = \partial S$ are obstructions to the null-sliceness for the links $L^{\bar{\epsilon}}$. This in turn gives a close relation between the Milnor invariants of the links $L^{\bar{\epsilon}}$ and the Vassiliev invariants of K (Theorem 4.4). More precisely, we show that there exists a sequence of natural numbers $\{l(n)\}_{n \in \mathbb{N}}$, with $l(n) > \log_2(\frac{n-5}{144})$, such that the following is true:

Theorem 1. *Let the notation be as above. Assume that $L^{\bar{\epsilon}}$ is n -slice. Then, all the Vassiliev invariants of orders $\leq l(2n-1)$ vanish for K .*

The sequence $\{l(n)\}_{n \in \mathbb{N}}$ is defined in §3. A key idea we will introduce is to define n -unknotted Seifert surfaces. Roughly speaking, these are surfaces whose complement looks, modulo certain terms of the lower central series of its fundamental group, “simple” (e.g. like the complement of a Seifert surface of a trivial knot). A knot bounding such a surface is called n -unknotted. Our main result in this paper shows that certain Vassiliev invariants of a knot obstruct to n -unknottedness. (For details see Theorem 5.6.) A particularly interesting class of surfaces that we study, that is used to prove Theorem 1 above, is that of n -hyperbolic. Roughly speaking, a regular Seifert surface S is called n -hyperbolic, for some $n \in \mathbb{N}$, if its complement looks, modulo the first $(n+1)$ terms of its fundamental group, like the complement of a disc. These surfaces are introduced in §3. We prove the following:

Theorem 2. *If K is n -hyperbolic, then all the Vassiliev invariants of orders $\leq l(n)$ of K vanish. In particular, if K is n -hyperbolic for all $n \in \mathbb{N}$ then all the Vassiliev invariants of K vanish.*

To motivate our notion of n -hyperbolicity, let us recall that a link is n -slice if its components bound disjoint smooth surfaces V_i in the 4-ball B^4 and such that each curve on V , if pushed off into the complement, lies in the n -th term of the lower central series of $\pi_1(B^4 \setminus \cup V_i)$. The results in [Co] and [O] and in [L1] motivated our definition above. Our terminology though, is motivated by the form of the Seifert matrix of the knot. See Remark 5.3.

In [Gu], Gusarov studied knots with trivial invariants of bounded orders by means of combinatorics on knot projections. His notion of n -triviality relates a knot with vanishing finite type invariants to the unknot, via crossing changes on

knot diagrams. This process has to take into account all the intermediate knots that the knot passes through. Our notion on the other hand, involves only the knot in question and its complement, and it is, thus, closer to the spirit of classical geometric topology.

An interesting problem, arising from this work, is the question of whether the notion of n -hyperbolicity provides a complete geometric characterization of n -trivial knots. We conjecture that this is the case. More precisely we have the following:

Conjecture 3. *A knot K is n -trivial for all $n \in \mathbb{N}$ if and only if it is n -hyperbolic for all $n \in \mathbb{N}$.*

The reader is referred to §6 for further discussion.

Let us now describe in more detail the contents of the paper and some of the ideas that are involved in the proofs of the main results.

In §1 we recall basic facts about Vassiliev invariants and the results from [Gu] that we use in subsequent sections.

In §2 we study regular Seifert surfaces of knots. We introduce the notion of *good position* for bands in projections of Seifert surfaces. Let B be a band of a regular Seifert surface S , which is assumed to be in band-disc form, and let γ denote the core of B . Also, let γ^ϵ denote a push-off of γ . The main feature of a projection of S with respect to which B is in *good position* is the following: We may find a word W , in the free generators of $\pi = \pi_1(S^3 \setminus S)$, representing γ^ϵ and such that every letter in W is realized by a band crossing in the projection.

In §3 we introduce *n-hyperbolic* regular Seifert surfaces and we prove Theorem 2. The special projections of §2 allow us to connect Gusarov's notion of *n-triviality* to an algebraic n -triviality in π , and exhibit a correspondence between geometry in $S^3 \setminus S$ and algebra in π . Let us explain this in some more detail. By [Gu], to prove Theorem 2 it will be enough to show that an *n-hyperbolic* knot has to be $l(n)$ -trivial. Showing that a knot is *k-trivial* amounts to showing that it can be unknotted in $2^{k+1} - 1$ ways by changing crossings in a fixed projection. See Definition 1.2. Having the projections of §2 at hand, the main step in the proof of Theorem 2 becomes showing the following: If γ is the core of a band B in *good position* and $\gamma^\epsilon \in \pi^{(m+1)}$, then we can trivialize B in $2^{l(m)+1} - 1$ ways (for the precise statement see Proposition 3.3). Here $\pi^{(m+1)}$ denotes the $(m + 1)$ -th term

of the lower central series of π .

The proof of Proposition 3.3 is based on a careful analysis of the geometric combinatorics of projections of the sub-arcs of γ representing simple commutators. We show that eventually γ may be decomposed into a disjoint union of “nice” arcs for which the desired conclusion follows by Dehn’s Lemma.

In §4 we start by recalling the Definition of n -slice and a few results from [Co] and [O]. Then we show how Theorem 1 follows from Theorem 2.

In §5 we introduce n -unknotted Seifert surfaces and we prove various generalizations Theorem 2. See Theorems 5.4 and 5.6.

Finally, in §6 we discuss some consequences and state some questions that arise from this work.

Acknowledgment. We thank James Conant for a careful reading of this paper. His examples (see Remarks 3.3* and 3.14*) were crucial to us in clarifying the subtleties of the layouts of commutators. Also, we thank the referee for his/her comments that were very helpful in improving the exposition of the manuscript.

1. VASSILIEV KNOT INVARIANTS AND GUSAROV’S n - TRIVIALITY

A *singular knot* $K \subset S^3$ is an immersed curve whose only singularities are finitely many transverse double points.

Let $\mathcal{K}_n$ be the rational vector space generated by the set of ambient isotopy classes of oriented, singular knots with exactly n double points. In particular $\mathcal{K} = \mathcal{K}_0$ is the space generated by the set of isotopy classes of oriented knots.

A knot invariant V can be extended to an invariant of singular knots by defining

$$\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \diagup \quad \diagdown \end{array} - \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}$$

for every triple of singular knots which differ at one crossing as indicated. In particular, $\mathcal{K}_n$ can be viewed as a subspace of $\mathcal{K}$ for every n , by identifying any singular knot in $\mathcal{K}_n$ with the alternating sum of the 2^n knots obtained by resolving its double points. Hence, we have a subspace filtration

$$\dots \subset \mathcal{K}_n \dots \subset \mathcal{K}_2 \subset \mathcal{K}_1 \subset \mathcal{K}$$

Definition 1.1. A Vassiliev knot invariant of order $\leq n$ is a linear functional on the space $\mathcal{K}/\mathcal{K}_{n+1}$. The invariants of order $\leq n$ form a subspace $\mathcal{V}_n$ of $\mathcal{K}^*$, the annihilator of the subspace $\mathcal{K}_{n+1} \subset \mathcal{K}$. We will say that an invariant v is of order n if v lies in $\mathcal{V}_n$ but not in $\mathcal{V}_{n-1}$.

Clearly, we have a filtration

$$\mathcal{V}_0 \subset \mathcal{V}_1 \subset \mathcal{V}_2 \subset \dots$$

To continue we need to introduce some notation. Let $D = D(K)$ be a diagram of a knot K , and let $\mathcal{C} = \mathcal{C}(D) = \{C_1, \dots, C_m\}$ be a collection of disjoint non-empty sets of crossings of D . Let us denote by $2^{\mathcal{C}}$ the set of all subsets of $\mathcal{C}$. Finally, for an element $C \in 2^{\mathcal{C}}$ we will denote by D_C the knot diagram obtained from D by switching the crossings in all sets contained in C . So, all together, we can get 2^m different knot diagrams from the pair $(D, \mathcal{C})$. Notice that each $C_i \in \mathcal{C}$ may contain more than one crossings.

Definition 1.2. ([Gu]) Two knots K_1 and K_2 are called *n-equivalent*, if K_1 has a knot diagram D with the following property: There exists $\mathcal{C} = \{C_1, \dots, C_{n+1}\}$, a collection of $n+1$ disjoint non-empty sets of crossings of D , such that D_C is a diagram of K_2 for every non empty $C \in 2^{\mathcal{C}}$. A knot K which is *n-equivalent* to the trivial knot will be called *n-trivial*.

By [Ya] every knot is *1-trivial*. The following theorem will be useful to us in the subsequent sections.

Theorem 1.3. ([Gu], [N-S]) Two knots K_1 and K_2 are *n-equivalent* if and only if all of their Vassiliev invariants of order $\leq n$ are equal. In particular, a knot K is *n-trivial* if and only if all its Vassiliev invariants of order $\leq n$ vanish.

2. REGULAR SEIFERT SURFACES OF A KNOT

2a. Generalities

Let K be an oriented knot in S^3 . A *Seifert surface* of K is an oriented, compact, connected, bicollared surface S , embedded in S^3 such that $\partial S = K$.

A *spine* of S is a bouquet of circles $\Sigma \subset S$, which is a deformation retract of S .

Definition 2.1. A Seifert surface S of a knot K is called *regular* if it has a spine Σ whose embedding in S^3 , induced by the embedding $S \subset S^3$, is isotopic to the standard embedding of a bouquet of circles. We will say that Σ is a *regular spine* of S .

Let $\Sigma_n \subset S^3$, be a bouquet of n circles based at a point p . A regular projection of Σ_n is a projection of Σ_n onto a plane with only transverse double points as possible singularities. Starting from a regular projection of Σ_n , we can construct an embedded compact oriented surface as follows: On the projection plane, let D^2 be a disc neighborhood of the basepoint p , which contains no singular points of the projection. Then, D^2 intersects the projection of Σ_n in a bouquet of $2n$ arcs and there are n arcs outside D^2 . We first replace each of the arcs outside D^2 by a flat band with the original arc as its core. Here a band being flat means we have an immersion when the band is projected onto the plane. That is to say that the only singularities the band projection has are these at the double points of the original arc projection so that bands overlap themselves exactly when the arcs overcross themselves.

Let S denote the surface obtained by the union of the disc D^2 and these flat bands, to which some full twists are added if necessary. We say that S is a surface *associated* to the given regular plane projection of Σ_n . We will also say that the surface S is in a *disc-band form*. A *band crossing* of S is obviously defined, and they are in one-one correspondence with crossings on the regular plane projection of Σ_n . We certainly have the freedom to move the full twists added to the bands anywhere. So we assume that all the twists of the band are moved near the ends of the bands. We may sometimes abuse the notation by not distinguishing a band and its core and only take care of the twists at the end of an argument.

Now let S be a regular surface of genus g . Pick a basepoint $p \in S$, and let Σ_n , $n = 2g$, be a regular spine of S such that p is the point on Σ_n where all circles in Σ_n meet. Let $\gamma_1, \beta_1, \dots, \gamma_g, \beta_g$ be the circles in Σ_n oriented so that they form a symplectic basis of $H_1(S)$. Assume further that a disc neighborhood of p in S is chosen so that its intersection with Σ_n consists of $2n$ arcs.

Lemma 2.2. Let S be a regular Seifert surface, with Σ_n a regular spine. The embedding $\Sigma_n \subset S^3$ has a regular plane projection as shown in Figure 1 below, where b is a braid of index $2n$, such that the regular Seifert surface S is isotopic

to a surface associated to that projection of Σ_n .

Proof. Let W_n be a bouquet of n circles, all based at a common point q . Then, Σ_n induces an embedding of W_n in S^3 . Let us begin with a regular plane projection of W_n , such that in a neighborhood D of q in the projection plane, the $2n$ arcs in $D \cap W_n$ are ordered and oriented in the same way as the arcs of Σ_n in the chosen disc neighborhood of p in S . Then, after a possible adjustment by adding some small kinks, S is isotopic to the surface associated to this projection. Since Σ_n is isotopic to the standard embedding of W_n in S^3 , we may switch the arcs in D , so that the arcs of Σ_n outside D are isotopic to the standard embedding. We may then record these switches by the braid b . $\square$




Figure 1. A projection of a regular spine of a Seifert surface

2b. Good position of a band in a regular spine

Let us consider $\mathbb{R}^3 \subset S^3$ and a decomposition $\mathbb{R}^3 = \mathbb{R} \times \mathbb{R}^2$, and take the factor $\mathbb{R}^2$ as a fixed projection plane P from now on. Also, we will fix a coordinate decomposition (t, s) of P . Now let l denote the t -axis on P , and let $H_+ = \{(t, s) \in P \mid s > 0\}$ and $H_- = \{(t, s) \in P \mid s < 0\}$.

To continue assume that S is a regular Seifert surface and fix a projection, $p : S \longrightarrow P$ in the disc-band form. Assume that the bands (their cores) of S are all transverse to l . Let B be a band of S , and let γ be the core of B . By a *sub-band* B' of B , we mean a band on B whose core γ' is a sub-arc of γ .

Definition 2.3. We will say that a band B is in *good position* with respect to the projection iff the following conditions hold:

- a) The band B is flat.
- b) For every band $A \neq B$, the intersection $H_+ \cap A$ consists of a single sub-band with no self-crossings, and these sub-bands are all disjoint.

- c) All the self-crossings of B occur in H_+ , and are all undercrossings (resp. overcrossings). Moreover, the intersection $H_+ \cap B$ consists of finitely many sub-bands $B_0, B_1, \dots, B_k$ such that
- i) they have no self-crossings;
 - ii) the B_j 's, $j \neq 0$, are disjoint with each other, and each crosses exactly once under (resp. over) B_0 , or one of the sub-bands in b).
- d) The crossings between B and any other band that occur in H_- are all overcrossings (resp. undercrossings).

An example of a projection as described in Definition 2.3 is shown in Figure 6, at the end of this section.

To continue, let S be a regular Seifert surface and fix a projection as described in Lemma 2.2. Let g be the genus and let $A_1, B_1, \dots, A_g, B_g$ denote the bands of S . Moreover, let $\gamma_1, \beta_1, \dots, \gamma_g, \beta_g$ denote the cores of $A_1, B_1, \dots, A_g, B_g$, respectively. We orient the core curves so that they give a symplectic basis of $H_1(S)$. Finally, let $x_1, y_1, \dots, x_g, y_g$ be small linking circles of the bands such that

i) $\text{lk}(x_i, \gamma_j) = \text{lk}(y_i, \beta_j) = \delta_{ij}$;

ii) $\text{lk}(y_i, \gamma_j) = \text{lk}(x_i, \beta_j) = 0$ and

iii) their projections on the plane P are simple curves disjoint from each other.

Clearly, $x_1, y_1, \dots, x_g, y_g$ represent free generators of $\pi_1(S^3 \setminus S)$.

Lemma 2.4. *For every band B of S , there exists a projection of S with respect to which B is in good position.*

Proof. Let us start with the projection fixed before the statement of the Lemma, and let l and H_+, H_- be as before Definition 2.3.

Let $\alpha_1, \hat{\alpha}_1, \dots, \alpha_{2g-1}, \hat{\alpha}_{2g}$ denote the hooks in Figure 1 on the top of the braid b . They are sub-bands of $A_1, B_1, \dots, A_g, B_g$, respectively. We move the projection of S so that l intersects each of the hooks at exactly two points and we have that the intersection $H_+ \cap p(S)$ is equal to $\alpha_1 \cup \hat{\alpha}_1 \cup \dots \cup \alpha_{2g-1} \cup \hat{\alpha}_{2g}$. Thus the entire braid b is left below l , in H_- . To continue, we choose another horizontal line l_0 below l , so that b lies between l_0 and l , and only the disc part of the surface is left below l_0 . Finally, we draw more horizontal lines $l_1, l_2, \dots, l_m = l$ such that the braid b has exactly one crossing between l_{i-1} and l_i for $i = 1, 2, \dots, m$.

Without loss of generality we may assume that $B = A_1$.

Observe that since b is a braid, each band crossing of A_1 under some band A can be slid all the way up, by using the *finger moves* of Figure 2. That is, we can slide a short sub-band of A_1 , which is underneath A at the crossing, up following A until it becomes a small hook above l undercrossing the hook of A .

To isotope the band A_1 into good position, we start with the lowest undercrossing of A_1 under, say some band A , between l_{i-1} and l_i , for some i . We slide it up above l and still call the resulting band A_1 . Now between l_i and l_{i+1} , if there is an undercrossing of A in the original picture, we will have two new undercrossings of the modified A_1 . We slide these two new undercrossings of A_1 up, above l , along the same way as we slide the undercrossing of A between l_i and l_{i+1} up above l .




Figure 2. Sliding an undercrossing across a band

Since b has only finitely many crossings, this procedure will slide all undercrossings of A_1 up above l , to make A_1 in good position.




Figure 3. Twists on a band realized as kinks and nested kinks

The condition a) of Definition 2.3 can also be satisfied by further isotopy which first moves the twists on the band B to a place around the line l and then changes them to a family of “nested kinks”. See Figure 3. $\square$

Remark 2.5. Notice that a similar procedure can be carried out for B with overcrossings replacing undercrossings and vice versa.

To continue let K be a knot, and let S be a genus g Seifert surface of K . Let $N \cong S \times (-1, 1)$ be a bicollar of S in S^3 , such that $S \cong S \times \{0\}$, and let $N^+ = S \times (0, 1)$ (resp. $N^- = S \times (-1, 0)$). For a simple closed curve $\gamma \subset S$, denote $\gamma^+ = \gamma \times \{1/2\} \subset N^+$ and $\gamma^- = \gamma \times \{-1/2\} \subset N^-$.

Now let the projection $p : S \longrightarrow P$, on the (t, s) -plane be as in Lemma 2.2. We denote by $z_1, z_2, \dots, z_s$ the generators of $\pi_1(S^3 \setminus S)$ arising from the Wirtinger presentation associated to the fixed projection of S (see for example [Ro]). The generators z_i are in one to one correspondence with the arcs of the projection between two consecutive undercrossings. Moreover, every z_i is a conjugate of one of the free generators fixed earlier.

In Figure 2 we have indicated the Wirtinger generators by small arrows under the bands. The directions of the arrows are determined by the fixed orientations of the free generators $x_1, y_1, \dots, x_g, y_g$. Notice that these free generators can be chosen as a part of the Wirtinger generators corresponding to the hooks $\alpha_1, \hat{\alpha}_1, \dots, \alpha_{2g-1}, \hat{\alpha}_{2g}$, respectively.

Lemma 2.6. (The geometric rewriting) *Let $\gamma \subset S$ be a simple closed curve represented by the core of a band B of S , and let γ^ϵ ($\epsilon = \pm$) be one of the push off's of γ . Moreover, let $W = z_{i_1} \cdots z_{i_m}$ be a word representing γ^ϵ in terms of the Wirtinger generators of the projection, and let $W' = W'(x_i, y_j)$ be the word obtained from W by expressing each z_{i_r} in terms of the free generators. Then, there exists a projection*

$$p' : S \longrightarrow P$$

which is obtained from p by isotopy, and such that every letter in W' is realized by an undercrossing of B with one of the hooks above l .

Proof. Suppose that $\epsilon = +$. Assume first that B is flat. Then every z_{i_r} in W can be realized by an undercrossing of B with another band or itself. We claim that the projection obtained in Lemma 2.4 has the desired properties.

To see that, let us begin with three Wirtinger generators z_i, z_j, z_k , around a crossing of the projection as show in Figure 2 (a). Then we have $z_k = z_i z_j z_i^{-1}$. Notice that γ^+ , which is drawn by the dotted arrow, picks up z_k at the undercrossing in (a). After performing a *finger move* in (b), each of z_i, z_j, z_i^{-1} is realized by an undercrossing. Thus the geometric equivalent of replacing each z_{i_j} , in $W = z_{i_1} \cdots z_{i_m}$, by its expression in terms of the free generators is to slide

an undercrossing of B until it reaches the appropriate hook. Now the desired conclusion follows easily from this observation.

If B is not flat, we first assume that the twists are all near one of the ends of B . Let the Wirtinger generator near that end of B be z_0 . Then $W = z_0^k z_{i_1} \cdots z_{i_m}$, and every z_{i_r} can be realized by an undercrossing of B with another band or itself. The previous argument still works in this case for each z_{i_j} . For z_0^k , we may first move the twists up to a place around l and then replace the twists by a family of nested kinks, like in the last part of the proof of Lemma 2.4. This finishes the proof of Lemma 2.6. $\square$

2c. Lower central series and curves on surfaces

For a group G let $[G, G]$ denote the commutator subgroup of G . The lower central series $\{G^{(m)}\}_{m \in \mathbb{N}}$ of G is defined by $G^{(1)} = G$ and

$$G^{(m+1)} = [G^{(m)}, G]$$

for $m \geq 1$.

We begin by recalling some commutator identities that will be useful to us later on. See [KMS].

Proposition 2.7. (Witt-Hall identities) *Let G be a group and let k, m and l be positive integers. Suppose that $x \in G^{(k)}$, $y \in G^{(m)}$ and $z \in G^{(l)}$. Then*

- a) $[G^{(k)}, G^{(m)}] \subset G^{(k+m)}$ or $xy \equiv yx \pmod{G^{(k+m)}}$
- b) $[x, zy] = [x, z] [x, y] [[y, x], z]$
- c) $[xy, z] = [y, z] [[z, y], x] [x, z]$
- d) $[x, [y, z]] [y, [z, x]] [z, [x, y]] \equiv 1 \pmod{G^{(k+l+m+1)}}$
- e) If $g \equiv g' \pmod{G^{(k)}}$ and $y \in G^{(m)}$ then $[g, y] \equiv [g', y] \pmod{G^{(k+m)}}$ and $[y, g] \equiv [y, g'] \pmod{G^{(k+m)}}$.

Let F be a free group of finite rank and let $\mathcal{A} = \{a_1, \dots, a_k\}$ be a set of (not necessarily free) generators of F . Let a be an element in F and let $W = W_a(\mathcal{A})$ be a word in $a_1, \dots, a_k$ representing a . Think of W as given as a list of spots in which we may deposit letters $a_1^{\pm 1}, a_2^{\pm 1}, \dots, a_k^{\pm 1}$. Now let $\mathcal{C} = \mathcal{C}(W) = \{C_1, \dots, C_m\}$ be a collection of disjoint non-empty sets of spots (or letters) in W . Let us denote by $2^{\mathcal{C}}$ the set of all subsets of $\mathcal{C}$. Finally, for an element $C \in 2^{\mathcal{C}}$ we will denote by

W_C the word obtained from W by substituting the letters in all sets contained in C by 1.

Definition 2.8. ([N-S]) *The element $a \in F$ is called n -trivial, with respect to $\mathcal{A}$, if it has a word presentation $W = W_a(\mathcal{A})$ with the following property: There exist a collection of $n + 1$ disjoint non-empty sets of letters, say $\mathcal{C} = \{C_1, \dots, C_{n+1}\}$, in W such that W_C represents the trivial element for every non-empty $C \in 2^{\mathcal{C}}$.*

We will say that $a \in F$ is n -trivial if it is n -trivial with respect to a set of generators. The following lemma shows that this definition depends neither on the word presentation nor the set of generators used.

Lemma 2.9. *If $a \in F$ is n -trivial with respect to a generating set $\mathcal{A}$, then it is n -trivial with respect to every generating set of F .*

Proof. Let $W = W_a(\mathcal{A}) = a_{i_1} a_{i_2} \dots a_{i_s}$ be a word for a satisfying the properties in Definition 2.8 and let $\mathcal{A}'$ be another set of generators for F . By expressing each a_{i_j} as a word of elements in $\mathcal{A}'$ we obtain a word $W' = W'_a(\mathcal{A}')$ which satisfies the requirements of n -triviality with respect to $\mathcal{A}'$. $\square$

Lemma 2.10. *If a lies in $F^{(n+1)}$, then it is n -trivial.*

Proof. Observe that a basic commutator $[a, b] = aba^{-1}b^{-1}$ is 1-trivial by using $\mathcal{C} = \{\{a, a^{-1}\}, \{b, b^{-1}\}\}$, and induct on n . $\square$

Clearly, we do not change the n -triviality of a word by inserting a *canceling pair* xx^{-1} or $x^{-1}x$, where x is a generator. We will use the following definition to simplify the exposition.

Definition 2.11. *A simple commutator is a word in the form of $[A, x^{\pm 1}]$ or $[x^{\pm 1}, A]$ where x is a generator and A is a simple commutator of shorter length. A simple quasi-commutator is a word obtained from a simple commutator by finitely many insertions of canceling pairs.*

By Proposition 2.7, any word representing an element in $F^{(n)}$ can be changed to a product of simple quasi-commutators of length $\geq n$ by finitely many insertions of canceling pairs. A simple quasi-commutator of length $> n$ is clearly n -trivial.

To continue, let S be a regular Seifert surface of a knot K . For a loop $\alpha \subset S^3 \setminus S$, we will denote by $[\alpha]$ its homotopy class in $\pi = \pi_1(S^3 \setminus S)$.

Suppose that $S, \gamma, B, p' : S \longrightarrow P$ and $[\gamma^\epsilon] = W'(x_i, y_i) = W'$ are as in the statement of Lemma 2.6. Suppose δ is a sub-band of γ and $[\delta^\epsilon] = W''$ is a sub-word of W' . Assume that W'' represents an element in $\pi^{(n)}$.

Lemma 2.12. (The geometric realization) *There exists a projection $p_1 : S \rightarrow P$ with the following properties:*

- i) $p_1(S)$ is obtained from $p'(S)$ by a finite sequence of band Reidemeister moves of type II;
- ii) B is in good position with respect to the new projection;
- iii) the word $W^* = W^*(x_i, y_i)$ one reads out from δ (with respect to the new projection p'), by picking up one letter for each crossing of δ underneath the hooks, is a product of simple quasi-commutators of length $\geq n$.

Proof. Since W'' represents an element in $\pi^{(n)}$, we may change W'' to a product of simple quasi-commutators by finitely many insertions of cancelling pairs. Such an insertion of a cancelling pair can be realized geometrically by a finger move (type II Reidemeister move). We will create a region in the projection plane to perform such a finger move. This region is a horizontal long strip below the line l and its intersection with $p'(S)$ consists of vertical straight flat bands. See Figure 4.




Figure 4. Realizing an insertion of $x^{-1}x$ or xx^{-1} by isotopy

As shown in Figure 4, there are two situations corresponding to insertions of $x^{-1}x$ or xx^{-1} . In one of the cases, we shall either let the finger go over one of the vertical flat bands connected to the x -hook (in the case that the x -hook does not belong to B) or push that vertical flat band along with the finger move (in the case that the x -hook belongs to B). Furthermore, if some vertical flat bands belonging to B block the way of the finger move, we will make more insertions by pushing these vertical flat bands along with the finger move. Finally, with all these done, we may easily modify the projection further to make B still in good position and ready to do the next insertion. $\square$

2d. An example illustrating the good position of a band

In Figure 5, we show an example of a regular Seifert surface of genus one.




Figure 5. A 2-hyperbolic surface

The cores of the bands A and B of the surface S have been drawn by the dashed, oriented curves γ_A and γ_B , respectively. The fundamental group $\pi = \pi_1(S^3 \setminus S)$ is freely generated by x and y . We have $[\gamma_A^+] \in \pi^{(3)}$, where γ_A^+ is the push-off of γ_A along the positive normal vector of the surface S pointing upwards the projection plane. In fact, from the Wirtinger presentation obtained from the given projection we have $[\gamma_A^+] = [zw^{-1}] = [[x, y], y^{-1}]$. Such a surface will be called *2-hyperbolic* in Definition 3.1. Now we modify the projection of S , so that A is in good position. The resulting projection is shown in Figure 6. Here we have only drawn the cores of the bands. The solid (dashed, resp.) arc corresponds to the band A (B , resp.) of Figure 6. The word we read out when traveling along the solid arc, one letter for each crossing underneath the hooks, is exactly $W = [[x, y], y^{-1}]$. The reader can verify that the two crossings marked by “%” on

the left side of the y -hook and the two crossings marked by “*” can be used to trivialize the band A in three ways.




Figure 6. The band A in good position

Remark 2.13. There is an obvious collection $\mathcal{C}$ of three sets of letters of W , so that W becomes a trivial word whenever we delete letters in a non-empty $C \in 2^{\mathcal{C}}$ from W (see proof of Lemma 2.10). The projection of Figure 6 has the property that every letter in the word $W = [[x, y], y^{-1}]$ is realized by a band crossing. So we obtain a collection of sets of crossings, also denoted by $\mathcal{C}$. However, as the reader can verify, the image of γ_A on the surface S_C , obtained from S by switching the crossings in C , will not always be homotopically trivial in $S^3 \setminus S_C$.

3. COMMUTATOR HALF BASES AND VASSILIEV INVARIANTS

In this section we undertake the study of *regular* Seifert surfaces, whose complement looks, modulo the first $n + 1$ terms of the lower central series of its fundamental group, like the complement of a null-isotopy. Our main goal is to show that the existence of such a surface for a knot K forces its Vassiliev invariants of certain orders to vanish.

3a. Definitions and the main result

Before we are able to state our main result in this section we need some notation and terminology. Let K be a knot in S^3 and let S be a Seifert surface of K , of genus g . Throughout this paper a *basis* of S will be a collection of $2g$

non-separating simple closed curves $\{\gamma_1, \beta_1, \dots, \gamma_g, \beta_g\}$ that represent a symplectic basis of $H_1(S)$. That is we have $\mathcal{I}(\gamma_i, \gamma_j) = \mathcal{I}(\beta_i, \beta_j) = \mathcal{I}(\beta_i, \gamma_j) = 0$, for $i \neq j$, and $\mathcal{I}(\gamma_i, \beta_i) = 1$, where $\mathcal{I}$ denotes the intersection form on S . Each of the collections $\{\gamma_1, \dots, \gamma_g\}$ and $\{\beta_1, \dots, \beta_g\}$ will be called a *half basis*.

To continue let $\pi = \pi_1(S^3 \setminus S)$. For a basis $\mathcal{B} = \{\gamma_1, \beta_1, \dots, \gamma_g, \beta_g\}$ of $H_1(S)$ let $\mathcal{B}^* = \{x_1, y_1, \dots, x_g, y_g\}$ denote elements in π representing the dual basis of $H_1(S^3 \setminus S)$.

For a subset $\mathcal{A}$ of $\mathcal{B}$, let $\mathcal{G}_{\mathcal{A}}$ denote the normal subgroup of π generated by the subset of $\mathcal{B}^*$ corresponding to $\mathcal{A}$. Moreover, we will denote by $\pi_{\mathcal{A}}$ (resp. $\phi_{\mathcal{A}}$) the quotient $\pi/\mathcal{G}_{\mathcal{A}}$ (resp. the quotient homomorphism $\pi \longrightarrow \pi/\mathcal{G}_{\mathcal{A}}$). Finally, $\pi_{\mathcal{A}}^{(m)}$ will denote the m -th term of the lower central series of $\pi_{\mathcal{A}}$. For the following definition it is convenient to allow $\mathcal{A}$ to be the empty set and have $\pi_{\mathcal{A}} = \pi$.

Definition 3.1. Let $n \in \mathbb{N}$. A regular Seifert surface S is called *n-hyperbolic*, if it has a half basis $\mathcal{A}$ represented by circles in a regular spine Σ with the following property: There is an ordering, $\gamma_1, \dots, \gamma_g$, of the elements in $\mathcal{A}$ such that either $\phi_{\mathcal{A}_{i-1}}([\gamma_i^+])$ or $\phi_{\mathcal{A}_{i-1}}([\gamma_i^-])$ lies in $\pi_{\mathcal{A}_{i-1}}^{(n+1)}$. Here $\mathcal{A}_k = \{x_1, y_1, \dots, x_k, y_k\}$ for $k = 1, \dots, g$ and $\mathcal{A}_0$ is the empty set. The boundary of such a surface will be called an *n-hyperbolic knot*.

Remark. A particularly interesting case of Definition 3.1 is when the surface S contains a half basis $\gamma_1, \dots, \gamma_g$ such that $[\gamma_i^+]$ or $[\gamma_i^-]$ lie in $\pi^{(n+1)}$.

In order to state our main result in this section we need some notation. For $m \in \mathbb{N}$, let $q(m)$ be the quotient of division of m by six (that is $m = 6q(m) + r_1$, $0 \leq r_1 \leq 5$). Let the notation be as in Definition 3.1. For $i = 1, \dots, g$, let x_i denote the free generator of π that is dual to $[\gamma_i]$. Let l_i denote the number of distinct elements in $\{x_1, y_1, \dots, x_g, y_g\}$, that are different than x_i and whose images under $\phi_{\mathcal{A}_{i-1}}$ appear in a (reduced) word, say W_i , representing $\phi_{\mathcal{A}_i}([\gamma_i^+])$ or $\phi_{\mathcal{A}_i}([\gamma_i^-])$. Write W_i as a product, $W_i = W_i^1 \dots W_i^{s_i}$, of elements in $\pi_{\mathcal{A}_{i-1}}^{(n+1)}$ and partition the set $\{W_i^1, \dots, W_i^{s_i}\}$ into disjoint sets, say $\mathcal{W}_i^1, \dots, \mathcal{W}_i^{t_i}$ such that: i) $k_i^1 + \dots + k_i^{t_i} = l_i$, where k_i^j is the number of distinct elements in $\mathcal{A}_i$ involved in $\mathcal{W}_i^j$ and ii) for $a \neq b$, the sets of elements from $\mathcal{A}_i$ appearing $\mathcal{W}_i^a$ and $\mathcal{W}_i^b$ are disjoint. Let

$$k_i = \min\{k_i^1, \dots, k_i^{t_i}\}.$$

and let

$$q_{\gamma_i} = \begin{cases} q(n+1), & \text{if } n < 6k \\ k_i + \left\lceil \log_2 \frac{n+1-6k_i}{6} \right\rceil, & \text{if } n \geq 6k. \end{cases}$$

Notice that

$$q(n+1) > \frac{n+1}{6} - 1 = \frac{n-5}{6} > \log_2\left(\frac{n-5}{6}\right).$$

Also, since $ab \geq a+b$ if $a, b > 1$, we have

$$k_i + \log_2\left(\frac{n+1-6k_i}{6}\right) > \log_2 k_i + \log_2\left(\frac{n+1-6k_i}{6}\right) > \log_2\left(\frac{n+1}{36}\right).$$

Thus, for $n > 5$, we have

$$q_{\gamma_i} > \log_2\left(\frac{n-5}{72}\right).$$

We define $l(n)$ by

$$l(n, S) = \min\{q_{\gamma_1} - 1, \dots, q_{\gamma_g} - 1\},$$

and

$$l(n) = \min\{l(n, S) \mid S \text{ is } n\text{-hyperbolic}\}.$$

We can now state our main result in this section, which is:

Theorem 3.2. *If K is n -hyperbolic, for some $n \in \mathbb{N}$, then K is at least $l(n)$ -trivial. Thus, all the Vassiliev invariants of K of orders $\leq l(n)$ vanish.*

From our analysis above, we see that $l(n) > \log_2\left(\frac{n-5}{144}\right)$ and in particular $\lim_{n \rightarrow \infty} l(n) = \infty$. Thus, an immediate Corollary of Theorem 3.2 is:

Corollary 3.2*. *If K is n -hyperbolic, for all $n \in \mathbb{N}$, then all its Vassiliev invariants vanish.*

Assume that S is in disc-handle form as described in Lemma 2.2 and that the cores of the bands form a symplectic basis of $H_1(S)$. Moreover, assume that the curves $\gamma_1, \dots, \gamma_g$, of Definition 3.1 can be realized by half of these cores. Let $\beta_1, \dots, \beta_g$ denote the cores of the other half bands and let $D = D(K)$ denote the knot diagram of K , induced by our projection of the surface. Also we may assume that the dual basis $\{x_1, y_1, \dots, x_g, y_g\}$ is represented by free generators of π , as before the statement of Lemma 2.4.

To continue with our notation, let C be a collection of band crossings on the projection of S . We denote by S_C (resp. D_C) the Seifert surface (resp. knot diagram) obtained from S (resp. D) by switching all crossings in C , simultaneously. For a simple curve $\gamma \subset S$ (or an arc $\delta \subset \gamma$), we will denote by γ_C (or δ_C) the image of γ (or δ) on S_C .

Let γ be the core of a band B in good position and suppose that it is decomposed into a union of sub-arcs $\eta \cup \delta$ with disjoint interiors, such that the word, say W , represented by δ^+ (or δ^-) in $\pi = \pi_1(S^3 \setminus S)$ lies in $\pi^{(m+1)}$. Let x be the generator of π corresponding to B and let l denote the number of distinct free generators, different than x , appearing in W . Write W as a product, $W = W_1 \dots W_s$, of commutators in $\pi^{(n+1)}$ and partition the set $\{W_1, \dots, W_s\}$ into disjoint sets, say $\mathcal{W}_1, \dots, \mathcal{W}_t$ such that: i) $k_1 + \dots + k_t = l$, where k_j is the number of distinct generators involved in $\mathcal{W}_j$ and ii) for $a \neq b$, the sets of generators appearing $\mathcal{W}_a$ and $\mathcal{W}_b$ are disjoint. Let

$$k = \min\{k_1, \dots, k_t\}.$$

We define

$$q_\delta = \begin{cases} q(n+1), & \text{if } n < 6k \\ k + \left\lceil \log_2\left(\frac{n+1-6k}{6}\right) \right\rceil, & \text{if } n \geq 6k. \end{cases}$$

The proof of Theorem 3.2 will be seen to follow from the following Proposition.

Proposition 3.3. *Let γ be the core of a band B in good position and suppose that it is decomposed into a union of sub-arcs $\eta \cup \delta$, such that the word represented by δ^+ (or δ^-) in $\pi = \pi_1(S^3 \setminus S)$ lies in $\pi^{(m+1)}$. Suppose that the word represented by η^+ (or η^-) is the identity.*

Let K' be the boundary of the surface obtained from S by replacing the sub-band of B corresponding to δ with a straight flat ribbon segment δ^ , connecting the endpoints of δ and above (resp. below) the remaining diagram. Then K and K' are at least l_δ -equivalent, where $l_\delta = q_\delta - 1$.*

The proof of Proposition 3.3 will be divided into several steps, and occupies all of §3. Without loss of generality we will work with δ^+ and γ^+ . In the course of the proof we will see that we may choose the collection of sets of crossings $\mathcal{C}$, required in the definition of l_δ -equivalence, to be band crossings in a projection of S . Moreover, for every non-empty $C \in 2^{\mathcal{C}}$, δ_C will be shown to be isotopic to a

straight arc, say δ^* , as in the statement above. Here $2^{\mathcal{C}}$ is the set of all subsets of $\mathcal{C}$.

In the rest of this paragraph, let us assume Proposition 3.3 and show how Theorem 3.2 follows from it.

Proof of Theorem 3.2. The proof will be by induction on the genus g of the surface S . If $g = 0$ then K is the trivial knot and there is nothing to prove. For $i = 1, \dots, g$, let A_i denote the band of S whose core corresponds to γ_i , and let B_i be the dual band.

By Definition 3.1 we have a band A_1 , such that the core γ satisfies the assumption of Proposition 3.3. We may decompose γ into a union of sub-arcs $\eta \cup \delta$ with disjoint interiors such that the word represented by δ^+ (resp. η^+) in $\pi = \pi_1(S^3 \setminus S)$ lies in $\pi^{(n+1)}$ (resp. is the empty word). Let K' be a knot obtained from K by replacing the sub-band of B corresponding to δ with a straight flat ribbon segment δ^* , connecting the endpoints of δ and above the remaining diagram, and let S' be the corresponding surface obtained from S . We will also denote the core of δ^* by δ^* .

By Proposition 3.3, K and K' are l_δ -equivalent. One can see that K' is n -hyperbolic, and it bounds an n -hyperbolic surface of genus strictly less than g .

Obviously, there is a circle on S' with δ^* as a sub-arc which bounds a disk D in $S^3 \setminus S'$. A surgery on S' using D changes S' to S'' with $\partial S'' = K'$, and we conclude that S'' is an n -hyperbolic regular Seifert surface with genus $g - 1$. Thus, inductively, K' , and hence K , is at least $l(n)$ -trivial. $\square$

Remark 3.3*. Let us close this paragraph by remarking that with the notation as in Definition 3.1, if we assume that each $[\gamma_i^\epsilon]$, $i = 1, \dots, g$ and $\epsilon = +$ or $-$, lies in $\pi^{(2)}$, then $K = \partial S$ has the trivial Alexander polynomial. Thus, since the only Vassiliev invariant of order two comes from the Alexander polynomial, K is *2-trivial*. Moreover, examples found by J. Conant ([C1]) indicate that if we require the stronger condition that $[\gamma_i^\epsilon] \in \pi^{(n+1)}$ in Definition 3.1, then the lower bound in Theorem 3.2 should have linear growth on n . However, Conant also has examples ([C]) demonstrating that with the weaker definition of n -hyperbolicity used in this paper, the logarithmic growth in the lower bound of Proposition 3.3 is necessary. More precisely, he constructs a family $\{K_n\}_{n \in \mathbb{N}}$ of knots such that: K_n is $(2^{n+2} - 4)$ -hyperbolic but, as computer calculations indicate, the lower bound

in Proposition 3.3 should be $2n$. We will not pursue an improvement of the lower bound in Theorem 3.2 here since, in the light of Proposition 6.2 and Conjecture 6.3, we are mainly interested in Corollary 3.2*.

3b. Nice arcs and simple commutators

In this paragraph we begin the study of the geometric combinatorics of arcs in good position and prove a few auxiliary lemmas required for the proof of Proposition 3.3. At the same time we also describe our strategy of the proof of 3.3.

Throughout the rest of section three, we will adapt the convention that the endpoints of δ or of any subarc $\tilde{\delta} \subset \delta$ representing a word in $\pi^{(m+1)}$, lie on the line l associated to our fixed projection. Let $W = c_1 \dots c_r$ be a word expressing δ^+ as a product of simple (quasi-)commutators of length $m+1$, and let $p_1(S)$ be a projection of S , as in Lemma 2.12. Then, each letter in W is represented by a band crossing in the projection. Now, let $\mathcal{C} = \{C_1, \dots, C_{m+1}\}$ be disjoint sets of letters obtained by applying Lemma 2.10 to the word W , so that W becomes a trivial word whenever we delete letters in a non-empty $C \in 2^{\mathcal{C}}$ from W (the resulting word is denoted by W_C).

Let y be a free generator appearing in W . We will say that the letters $\{y, y^{-1}\}$ constitute a *canceling pair*, if there is some $C \in 2^{\mathcal{C}}$ such that the word W_C can be reduced to the identity, in the free group π , by a series of deletions in which y and y^{-1} cancel with each other.

Ideally, we would like to be able to say that for every $C \in 2^{\mathcal{C}}$ the arc δ_C (obtained from δ by switching all crossings corresponding to C) is isotopic in $S^3 \setminus S_C$ to a straight segment connecting the end points of δ and above the remaining diagram. As remarked in 2.13, though, this may not always be the case. In other words not all sets of letters $\mathcal{C}$, that come from Lemma 2.10, will be suitable for geometric *m-triviality*. This observation leads us to the following definition.

Definition 3.4. Let S , B and δ be as in the statement of 3.3 and let $\tilde{\delta} \subset \delta$ be a subarc that represents a word W in $\pi^{(m+1)}$. Furthermore let δ^* be an embedded segment connecting the endpoints of $\tilde{\delta}$ and such that $\partial\delta^* \subset l$ and the interior of δ^* lies above the projection of S on the projection plane.

1) We will say that $\tilde{\delta}$ is *quasi-nice* if there exists a segment δ^* as above and such that either the interiors of δ^* and $\tilde{\delta}$ are disjoint, or $\tilde{\delta} = \delta^*$ and δ^* is the hook of the band B . Furthermore, if the interiors of δ^* and $\tilde{\delta}$ are disjoint then δ^* should not separate any set of crossings corresponding to a canceling pair in W on any of the hooks of the projection.

2) Let δ be a quasi-nice arc, and let δ^* be as in 1). Moreover, let S' denote the

surface $S \cup n(\delta^*)$, where $n(\delta^*)$ is a flat ribbon neighborhood of δ^* . We will say that δ is k -nice, for some $k \leq m + 1$, if there exists a collection $\mathcal{C}$ of k disjoint sets of band crossings on $\tilde{\delta}$, such that for every non-empty $C \in 2^{\mathcal{C}}$, the loop $(\delta^* \cup \tilde{\delta}_C)^+$ is homotopically trivial in $S^3 \setminus S'_C$, where $S'_C = S_C \cup n(\delta^*)$. We will say that every $C \in 2^{\mathcal{C}}$ trivializes $\tilde{\delta}$ geometrically.




Figure 7. Nice arcs representing simple 2-commutators

Notice that the arc in the example on the left side of Figure 7 is both 2-nice and quasi-nice while the one on the right side is not. In fact, one can see that all embedded arcs in good position representing simple 2-commutators are 2-nice.

Lemma 3.5. *Let δ be a subarc of the core of a band B , in a projection of a regular surface S . Let δ^* be a straight segment, connecting the endpoints of δ and let S' denote the surface $S \cup n(\delta^*)$. Suppose that the loop $(\delta^* \cup \delta)^+$ is homotopically trivial in $S^3 \setminus S'$. Then $n(\delta)$ can be isotoped onto $n(\delta^*)$ in $S^3 \setminus S$ relative to the endpoints.*

Proof. Since $\delta^* \cup \delta \subset S'$ is an embedded loop, by Dehn's Lemma (see for example [He] or [Ro]) we conclude that it bounds an embedded disc in $S^3 \setminus S'$. Then the claim follows easily. $\square$

Corollary 3.6. *Let γ and δ as in the statement of Proposition 3.3. Assume that δ is an q_δ -nice arc. Then the conclusion of the Proposition is true for δ .*

Proof. It follows immediately from Definitions 1.2 and 3.4 and Lemma 3.5. $\square$

For the rest of this paragraph we will focus on projections of arcs, in *good position*, that represent simple quasi-commutators. We will analyze the geometric combinatorics of such projections. This analysis will be crucial, in the next paragraphs, in showing that an arc δ as in Proposition 3.3 is q_δ -nice.

Let δ_1 be a subarc of δ presenting a simple quasi-commutator of length m , say c . Moreover, let δ_2 be another subarc of δ presenting a simple quasi-commutator equivalent to c or c^{-1} . We may change the orientation of δ_2 if necessary so that it presents a simple quasi-commutator equivalent to c . Then we may speak of the initial (resp. terminal) point $p_{1,2}$ (resp. $q_{1,2}$) of $\delta_{1,2}$; recall these points all lie on the line l .

Definition 3.7. Let $\hat{\delta}_1$ (resp. $\hat{\delta}_2$) be the segment on l going from p_1 to p_2 (resp. q_1 to q_2). We say that δ_1 and δ_2 are *parallel* if the following are true: i) At most one hook has its end points on $\hat{\delta}_1$ or $\hat{\delta}_2$ and both of its end points can be on only one of $\hat{\delta}_{1,2}$; ii) If a hook has exactly one point on some $\hat{\delta}_j$, say on $\hat{\delta}_1$, then $\hat{\delta}_1$ doesn't intersect the interior of $\delta_{1,2}$. iii) We have either $\hat{\delta}_1 \cap \hat{\delta}_2 = \emptyset$ or $\hat{\delta}_1 \subset \hat{\delta}_2$; iv) If $\hat{\delta}_{1,2}$ are drawn disjoint and above the surface S , the diagram $\delta_1 \cup \hat{\delta}_1 \cup \delta_2 \cup \hat{\delta}_2$ can be changed to an embedding by type II Reidemeister moves.

The reader may use Figure 8 to understand Definition 3.7. It should not be hard to locate the arcs $\hat{\delta}_{1,2}$ in each case in Figure 8. In the first two pictures, the straight arcs $\hat{\delta}_{1,2}$ have no crossings with $\delta_{1,2}$. Crossings between $\hat{\delta}_{1,2}$ and $\delta_{1,2}$ removable by type II Reidemeister moves are allowed to accommodate the modification of $\delta_{1,2}$ in Lemma 2.12. For example, in the last two pictures of Figure 8 one of $\hat{\delta}_{1,2} \subset l$ intersects both of $\delta_{1,2}$.




Figure 8. Various kinds of parallel arcs

Lemma 3.8. Assume that the setting is as in the statement of Proposition 3.3. Let c_1 and c_2 be equivalent simple quasi-commutators presented by sub-arcs $\delta_{1,2}$

of δ respectively, and let y be one of the free generators associated to the hooks of our fixed projection. Moreover, assume that $\delta_{1,2}$ are parts of a subarc ζ of δ presenting a simple quasi-commutator $W = c_1 y c_2^{-1} y^{-1}$. Then δ_1 and δ_2 are parallel.

Proof. By abusing the notation, we denote $\delta_1 = \tau_1 x \mu_1 x^{-1}$ and $\delta_2 = \tau_2 x \mu_2 x^{-1}$ where $\tau_1, \tau_2, \mu_1^{-1}, \mu_2^{-1}$ all present equivalent simple quasi-commutators. Furthermore, $\zeta = \delta_1 y \delta_2^{-1} y^{-1}$. For a subarc ν , up to symmetries, there are four possible ways for both of its endpoints to reach a certain z -hook so that $z\nu z^{-1}$ is presented by an arc in good position. See Figure 9, where the arc ν may run through the z -hook. We will call the pair of undercrossings $\{z, z^{-1}\}$ a *canceling pair*. Now let us consider the relative positions of τ_1, τ_2, μ_1 and μ_2 . Inductively, τ_i and μ_i^{-1} are parallel, for $i = 1, 2$. If $x\mu_1 x^{-1}$ is of type (I) in Figure 9, since τ_1 and μ_1^{-1} are parallel, τ_1 has to go the way indicated in Figure 10 (a).




Figure 9. Types of arcs presenting $z\nu z^{-1}$

If $x\mu_2 x^{-1}$ is also of type (I), there are two cases to consider. One case is to have the canceling pairs $\{x, x^{-1}\}$ in $x\mu_1 x^{-1}$ and $x\mu_2 x^{-1}$ both going underneath the x -hook at the left side, and the other case is to have them going underneath the x -hook at different sides. In the first case, in order to read the same word from τ_1 and τ_2 as well as from μ_1 and μ_2 , $\tau_1 x \mu_1 x^{-1}$ and $\tau_2 x \mu_2 x^{-1}$ has to fit like

in Figure 10 (b). This implies that δ_1 and δ_2 are parallel. In the second case (see Figure 10 (c)), in order that δ_i be parts of the arc $\zeta = \delta_1 y \delta_2^{-1} y^{-1}$, they have to go to reach the same y -hook.




Figure 10. The case when both $x\mu_1x^{-1}$ and $x\mu_2x^{-1}$ are of type (I)

But then we will not be able to read the same word through τ_1 and τ_2 . This shows that if $x\mu_1x^{-1}$ and $x\mu_2x^{-1}$ are both of type (I), δ_1 and δ_2 are parallel. There are many other cases which can be checked one by one in the same way as in Figure 10. The details are left to the patient reader. $\square$

So far we have been considering the projection of our surface on a plane P inside $\mathbb{R}^3 = P \times \mathbb{R}$. To continue, let us pass to the compactifications of $\mathbb{R}^3$ and P . We obtain a 2-sphere S_P^2 inside S^3 , and assume that our projection in Proposition 3.3 lies on S_P^2 . We may identify the image of l with the equator of S_P^2 , and the images of H_+ and H_- with the upper and lower hemisphere. We will interchange between P and S_P^2 whenever convenient.

Digression 3.9. Let δ , B be as in the statement of Proposition 3.3 and let x_0 denote the free generator of $\pi_1(S^3 \setminus S)$ corresponding to B . Suppose δ_1 and δ_2 are parallel subarcs of δ . and let $\hat{\delta}_{1,2}$ be as in Definition 3.7. We further assume

that the crossings between $\hat{\delta}_{1,2}$ and $\delta_{1,2}$ have been removed by isotopy. Let y be a free generator of $\pi = \pi_1(S^3 \setminus S)$. We assume that both δ_1 and δ_2 are sub-arcs of an arc $\zeta \subset \delta$ presenting $[c^{\pm 1}, y^{\pm 1}]$ (recall that $\delta_{1,2}$ present $c^{\pm 1}$). Then ζ is a union $\delta_1 \cup \tau_1 \cup \delta_2 \cup \tau_2$, where $\tau_{1,2}$ are segments each going once underneath the y -hook. One point of ζ is the same as one endpoint of one of $\delta_{1,2}$, say δ_2 . Let $\bar{\delta} := \delta \setminus (\delta_1 \cup \delta_2)$. By the properties of good position we see that in order for one of $\delta_{1,2}$, say δ_1 , not to be embedded on the projection plane it must run through the hook part of B , and the word representing δ_1 must involve x_0 . Moreover, good position imposes a set of restrictions on the relative positions of $\delta_{1,2}$ and the various subarcs of $\bar{\delta}$. Below we summarize the main features of the relative positions of $\delta_{1,2}$ and the various subarcs of $\bar{\delta}$; these features will be useful to us in the rest of the paper. We will mainly focus on the case that $\delta_{1,2}$ are embedded; the case of non-embedded arcs is briefly discussed in part b) of this Digression.

a) Suppose that $\delta_{1,2}$ are embedded on the projection plane P . Then the loop $\delta_1 \cup \hat{\delta}_2 \cup \delta_2 \cup \hat{\delta}_1$ separates S_P^2 into two discs, D_1 and D_2 . The intersections $D_{1,2} \cap \bar{\delta}$ consist of finitely many arcs. With the exception of at most one these arcs are embedded. One can see (see the two pictures on the left side of Figure 8) that the interiors of $\tau_{1,2}$ are disjoint from that of exactly one of $D_{1,2}$, say D_1 , and they lie in the interior of the other. We will call D_1 (resp. D_2) the *finite* (resp. *infinite*) disc corresponding to the pair $\delta_{1,2}$. Using the properties of good position one can see that for each component θ of $D_1 \cap \bar{\delta}$ one of the following is true:

(a₁) Both the endpoints of θ lie on $\hat{\delta}_2$ and θ can be pushed in the infinite disc D_2 after isotopy, or it represents a word w , such that the following is true: None of the letters appearing in the reduced form of w appears in the underlying commutator of c . Moreover for each free generator x appearing in w , $C^{\pm 1}$ contains inserted pairs $x^{\pm 1}x^{\mp 1}$. To see these claims, first notice that if one of the generators, say z , appears in the underlying commutator of c then the intersection of D_1 and the z -hook consists of two (not necessarily disjoint) arcs, say $\theta_{1,2}$, such that one point of $\partial(\theta_{1,2})$ is on δ_1 and the other on δ_2 . Moreover, both the endpoints of the z -hook lie outside D_1 in the infinite disc. Now a subarc of $\bar{\delta}$ in D_1 has the choice of either hooking with z in exactly the same fashion as $\delta_{1,2}$, or “push” $\theta_{1,2}$ by a finger move as indicated in Figure 4, and hook with some $x \neq z^{\pm 1}$. In order for the second possibility to occur, at least one of the endpoints of the x -hook must lie inside D ; by our discussion above this will not happen if x has already appeared in the

underlying commutator of c . The rest of the claim follows from the fact that the “top” of the x -hook must lie outside D_1

(a_2) One endpoint of θ is on $\hat{\delta}_1$ and the other on $\hat{\delta}_2$. Moreover, θ either represents $c^{\pm 1}$ or the trivial word or it represents a w , as in case (a_1) above.

(a_3) θ is a subarc of the hook corresponding to the band B , and it has one endpoint on δ_1 and the other on δ_2 or one point on δ_i and the other on $\hat{\delta}_j$ ($i, j = 1, 2$). Furthermore, if θ has one endpoint on δ_i and the other on $\hat{\delta}_j$ then i) the underlying commutator of c does not involve $x_0^{\pm 1}$; and ii) both endpoints of the x_0 -hook lie inside D_1 .

(a_4) One of $\delta_{1,2}$, say δ_1 , runs through the hook part of B and θ has one point on δ_1 and the other on $\hat{\delta}_j$ ($i, j = 1, 2$). Moreover, we have: i) The arc θ represents $w x_0^{\pm 1}$ where either $w = c^{\pm 1}$ or none of the letters appearing in there reduced form of w appears in the underlying commutator of c ; ii) the underlying commutator of c does not involve $x_0^{\pm 1}$; iii) if e is a simple quasi-commutator represented by a subarc $\tilde{\delta}$ such that $\delta_{1,2} \subset \tilde{\delta}$, then the underlying commutator of e does not involve $x_0^{\pm 1}$ (see also Lemma 3.12 a)).

b) Recall that x_0 is the free generator of π corresponding to B . Suppose that δ_1 is non-embedded. Then δ_1 to run through the hook of B and the word representing δ_1 must involve x_0 .

(b_1) It follows from the properties of good position that any subarc of $\theta \subset \delta$ that has its endpoints on different $\hat{\delta}_i$ has to represent $c^{\pm 1}$.

Lemma 3.10. *Let the setting be again as in the statement of Proposition 3.3, and let δ_1 be a subarc of δ representing a simple (quasi-)commutator. Moreover let h_0 denote the hook part of the band B . We can connect the endpoints of δ_1 by an arc δ_1^* , which is embedded on the projection plane and such that: i) δ_1^* lies on the top of the projection $p_1(S)$; ii) the boundary $\partial(\delta_1^*)$ lies on the line l ; and iii) either $\delta_1^* = h_0$ or the interiors of δ_1^* and δ_1 are disjoint and δ_1^* goes over at most one hook at most once.*

Proof. Suppose that δ_1 represents $W = [c, y^{\pm 1}]$ and let $\delta_1^{1,2}$ be the subarcs of δ_1 representing $c^{\pm 1}$ in W . By Lemma 3.8 $\delta_1^{1,2}$ are parallel; let $\hat{\delta}_1^{1,2}$ be the arcs of Definition 3.7 connecting the endpoints of $\delta_1^{1,2}$. Recall that there is at most one hook, say corresponding to a generator z , that can have its endpoints on $\delta_1^{1,2}$. If

$z = y$ then , using good position, we see that there is an arc δ_1^* as claimed above such that either $\delta_1^* = h_0$ or it intersects at most the y -hook in at most one point. If $z \neq y$ then we can find an arc α satisfying i) and ii) above and such that either $\alpha = h_0$ or α intersects the y -hook in at most one point and the intersections of α with the other hooks can be removed by isotopying α , relatively its endpoints. Thus the existence of δ_1^* follows again. $\square$

Definition 3.11. *An arc δ_1 representing a simple (quasi-)commutator will be called good if the arc δ^* of Lemma 3.10, connecting the endpoints of the δ_1 , doesn't separate any canceling pair of crossings in δ_1 .*

The reader can see that the arc in the picture of the left side of Figure 7 is good while the one on the right is not good.

The outline of the proof of 3.3. In 3.4 we defined the notions of quasi-niceness and k -niceness. By definition, a k -nice arc is quasi-nice. We will, in fact, show that the two notions are equivalent. More precisely, we show in Lemma 3.21 that a quasi-nice arc δ is q_δ -nice. This, in turn, implies Proposition 3.3. To see this last claim, notice that the arc δ in the statement of 3.3 is quasi-nice. Indeed, since the arc η in the statement of 3.3 represents the identity in π , good position and the convention about the endpoints of δ made in the beginning of 3b assure the following: Either the interior of η lies below l (and above the projection of $S \setminus n(\eta)$) and it is disjoint from that of δ or η is the hook part of B . In both cases we choose $\delta^* = \eta$.

To continue, notice that a good arc is by definition quasi-nice. The notion of a *good* arc is useful in organizing and studying the various simple quasi-commutator pieces of the arc δ in 3.3. In Lemma 3.15 we show that if an arc $\tilde{\delta}$ is good then it is $q_{\tilde{\delta}}$ -nice and in Lemma 3.19 we show that if $\tilde{\delta}$ is a product of good arcs then it is $q_{\tilde{\delta}}$ -nice. In both cases we exploit good position to estimate the number of “bad” crossings along $\tilde{\delta}$, that are suitable for algebraic triviality but may fail for geometric triviality. All these are done in 3c and 3d.

In 3e we begin with the observation that if $\tilde{\delta}$ is a product of arcs $\theta_1, \dots, \theta_s$ such that θ_i is q_{θ_i} -nice then $\tilde{\delta}$ is $q_{\tilde{\delta}}$ -nice (see Lemma 3.20). Finally, Lemma 3.21 is proven by induction on the number of “bad” subarcs that δ contains.

3c. Bad sets and good arcs

In this paragraph we continue our study of arcs in good position that represent simple quasi-commutators. Our goal, is to show that a good arc δ representing a simple quasi-commutator is q_δ -nice (see Lemma 3.15).

Let $W = [\dots[[y_1, y_2], y_3], \dots, y_{m+1}]$ be a simple (quasi-) commutator represented by an arc δ of the band B in good position. Suppose that the subarc of δ representing $W_1 = [\dots[[y_1, y_2], y_3], \dots, y_i]$, for some $i = 1, \dots, m+1$, runs through the hook part of B , at the stage that it realizes the crossings corresponding to y_i . The canceling pair corresponding to $\{y_i, y_i^{-1}\}$ will be called *the special canceling pair* of W .

Note: Let x_0 be the free generator of π corresponding to the band B . Notice that if the special canceling pair $\{y_i, y_i^{-1}\}$ is of type (III) or (IV) then we must have $y_i = x_0^{\pm 1}$; otherwise we must have $y_i \neq x_0^{\pm 1}$. Moreover, it follows by good position and Lemma 3.8 that if $\{y_i, y_i^{-1}\}$ is of type (I) or (II), then the arc representing $W_1 = [\dots[[y_1, y_2], y_3], \dots, y_{i-1}]$ is embedded; that is $y_j \neq x_0^{\pm 1}$ for $j = 1, \dots, i-1$.

Lemma 3.12. *Let $W = [\dots[[y_1, y_2], y_3], \dots, y_{m+1}]$ be a simple (quasi-) commutator represented by an arc δ of the band B in good position, and let x_0 be the free generator of π corresponding to the hook of B .*

- a) *Suppose that, for some $i = 1, \dots, m+1$, one of the canceling pairs $\{y_i, y_i^{-1}\}$ is the special canceling pair. Then, we have $y_j \neq x_0^{\pm 1}$ for all $i < j \leq m+1$.*
- b) *Let z be any free generator of π corresponding to one of the hooks of our projection. Then, at most two successive y_i 's can be equal to $z^{\pm 1}$.*

Proof. a) For $j > i$ let $c = [\dots[[y_1, y_2], y_3], \dots, y_{j-1}]$ and let $\delta_{1,2}$ be the arcs representing $c^{\pm 1}$ in $[c, y_j]$. By 3.8, $\delta_{1,2}$ are parallel. Let $\hat{\delta}_{1,2}$ be arcs satisfying Definition 3.7. Notice that the x_0 -hook can not have just one of its endpoints on $\hat{\delta}_{1,2}$. For, if the x_0 -hook had one endpoint on, say, $\hat{\delta}_1$, then $\hat{\delta}_1$ would intersect the interior of $\delta_{1,2}$. Now easy drawings will convince us that we can not form $W_1 = [c, x_0^{\pm 1}]$ without allowing the arc representing it to have self intersections below the line l associated to our projection. But this would violate the requirements of good position.

- b) By symmetry we may assume that $W = [\dots, [c^{\pm 1}, z^{\pm 1}], y_i, \dots, y_{m+1}]$, where c is a simple (quasi-)commutator of length $< m$. Let $\tilde{\delta}$ be the subarc of δ representing

$[c^{\pm 1}, z^{\pm 1}]$, and let $\delta_{1,2}$ be the subarcs of $\tilde{\delta}$ representing $c^{\pm 1}$. By Lemma 3.8, $\delta_{1,2}$ are parallel. Let $\hat{\delta}_{1,2}$ be as in Definition 3.7.

Without loss of generality we may assume that $z^{\pm 1}$ has already appeared in c . Thus the intersection $\delta_{1,2} \cap H_+$ is a collection of disjoint arcs $\{A_i\}$, each passing once under the z -hook, and with their endpoints on the line l . Let $A_{1,2}$ denote the innermost of the A_i 's corresponding to the two endpoints of the z -hook. Let $\alpha_{1,2}$ denote the segments of l connecting the endpoints of $A_{1,2}$, respectively. Observe that both of the endpoints of at least one of $\hat{\delta}_{1,2}$ must lie on α_1 or α_2 . There are three possibilities:

- (i) The endpoints of both $\hat{\delta}_{1,2}$ lie on the same $\alpha_{1,2}$, say on α_1 ;
- (ii) The endpoints of $\hat{\delta}_{1,2}$ lie on different $\alpha_{1,2}$;
- (iii) The endpoints of one of $\hat{\delta}_{1,2}$ lie outside the endpoints of $\alpha_{1,2}$.

Suppose we are in (i) above: Notice that both the endpoints of the arc $\tilde{\delta}$ must also lie on α_1 . By Definition 3.7 we see that both of the endpoints of any arc parallel to $\tilde{\delta}$ must also lie on α_1 . There are two possibilities for the relative positions of $\hat{\delta}_{1,2}$; namely $\hat{\delta}_1 \cap \hat{\delta}_2 = \emptyset$ or $\hat{\delta}_1 \subset \hat{\delta}_2$. Using Digression 3.9 we can see that in both cases we must have $y_i \neq z^{\pm 1}$.

If the last letter in $c^{\pm 1}$ was not $z^{\pm 1}$ then we are done. Suppose, now, that $c = [d, z^{\pm 1}]$. Then, A_1 must be part of one of $\delta_{1,2}$ and one endpoint of one of $\hat{\delta}_{1,2}$ must be an endpoint of α_1 . Moreover, observe that the last canceling pair in c can not be of type (I). By part a) this last canceling pair in c must be of type (II), and the only way that the last letter in d can be $z^{\pm 1}$ is to have $d = [e, z^{\pm 1}]$ where the last canceling pair is of type (I) or (II). Now easy drawings, using Digression 3.9, will convince us that this is not possible.

We now proceed with case (ii) above: Observe that the last letter in c can not be $z^{\pm 1}$; otherwise the endpoints of each of $\delta_{1,2}$ would be on the same $\alpha_{1,2}$. We first form $[c^{\pm 1}, z^{\pm 1}]$. The endpoints of the arc $\tilde{\delta}$ representing $[c^{\pm 1}, z^{\pm 1}]$ are now on the same $\alpha_{1,2}$, say on α_1 . Thus both of the endpoints of any arc parallel to $\tilde{\delta}$ must also lie on α_1 . Now we are in the situation of (i) above and the conclusion follows.

Finally, assume we are in case (iii) above: Notice that both endpoints of the arc $\tilde{\delta}$ representing $[c^{\pm 1}, z^{\pm 1}]$ lie outside $\alpha_{1,2}$. Thus $y_i \neq z^{\pm 1}$ or we are in the situation of (i) above. Using an argument similar to these of (i), (ii) we can see that we can't have the two last letters of $c^{\pm 1}$ being equal to $z^{\pm 1}$. $\square$

In order to continue we need some notation and terminology. We will write $W = [y_1, y_2, y_3, \dots, y_{m+1}]$ to denote the simple (quasi-)commutator

$$W = [\dots [[y_1, y_2], y_3], \dots, y_{m+1}].$$

Let $C_1, \dots, C_{m+1}$ be the sets of letters of Lemma 2.10, for W . Recall that for every $i = 1, \dots, m+1$ the only letter appearing in C_i is $y_i^{\pm 1}$.

Definition 3.13. We will say that the set C_i is bad if there is some $j \neq i$ such that i) we have $y_j = y_i = y$, for some free generator y ; and ii) the crossings on the y -hook, corresponding to a canceling pair $\{y_j, y_j^{-1}\}$ in C_j , are separated by crossings in C_i .

The problem with a bad C_i is that changing the crossings in C_i may not trivialize the arc δ geometrically.

For $i = 2, \dots, m+1$, let $c_i = [y_1, \dots, y_{i-1}]$ and let $\delta_{1,2}$ be the parallel arcs representing $c_i^{\pm 1}$ in $[c_i, y_i]$. Let $\bar{\delta} = \delta \setminus (\delta_1 \cup \delta_2)$. We will say that the canceling pair $\{y_i, y_i^{-1}\}$ is admissible if it is of type (I) or (II).

Note: If W doesn't involve the generator of π corresponding to the hook of B then every canceling pair in W is admissible.

Lemma 3.14. a) Let $W = [y_1, y_2, y_3, \dots, y_{m+1}]$ be a simple quasi-commutator represented by an arc δ in good position and let z be a free generator. Also, let $C_1, \dots, C_{m+1}$ be sets of letters as above. Suppose that C_i is bad and let $\{y_j, y_j^{-1}\}$ be a canceling pair in C_j , whose crossings on the z -hook are separated by crossings in C_i . Suppose, moreover, that the pair $\{y_i, y_i^{-1}\}$ is admissible. Then, with at most one exception, we have $j = i - 1$ or $j = i + 1$.

b) Let $w(z)$ be the number of the y_i 's in W that are equal to $z^{\pm 1}$. There can be at most $\left\lceil \frac{w(z)}{2} \right\rceil + 1$ bad sets involving $z^{\pm 1}$.

c) For every $j = 2, \dots, m+1$, at least one of $[y_1, \dots, y_{j-1}]$ and $[y_1, \dots, y_j]$ is represented by a good arc.

Proof. a) Let $c = [y_1, \dots, y_{i-1}]$, let $\delta_{1,2}$ be the parallel arcs representing $c^{\pm 1}$ in $[c, y_i]$ and let $\hat{\delta}_{1,2}$ be the arcs of Definition 3.7. Let C_z denote the canceling pair corresponding to $y_i^{\pm 1}$ in $[c, y_i]$. Moreover, let $\bar{\delta} = \delta \setminus (\delta_1 \cup \delta_2)$ and let $\bar{\delta}_c$ denote

the union of arcs in $\bar{\delta}$ such that i) each has one endpoint on $\hat{\delta}_1$ and one on $\hat{\delta}_2$ and ii) they do not represent copies of $c^{\pm 1}$ in W . By Lemma 3.8, and Digression 3.9, it follows that $\bar{\delta}_c = \emptyset$.

Without loss of generality we may assume that $j > i$. Also, we may, and will, assume that $y_{i+1} \neq z^{\pm 1}$.

First suppose that C_z is of type (II): By Lemma 3.12(a), it follows that if one of $\delta_{1,2}$ has runned through the hook part of B then $\bar{\delta} \cap \delta_{1,2} = \emptyset$. Thus the possibility discussed in (a_4) of Digression 3.9 doesn't occur. Now, by Lemma 3.8, it follows that in order for the crossings corresponding to C_z to separate crossings corresponding to a later appearance of $z^{\pm 1}$, we must have i) $y_j = z^{\pm 1}$ realized by a canceling pair of type (II) and ii) the crossings on the z -hook corresponding to y_j and y_j^{-1} lie below (closer to endpoints of the hook) these representing y_i and y_i^{-1} . But a moment's thought, using 3.9 and the assumptions made above, will convince us that in order for this to happen we must have $\bar{\delta}_c \neq \emptyset$; which is impossible.

Suppose C_z is of type (I): Up to symmetries, the configuration for the arc $\tilde{\delta}$, representing $[c, y_i]$, is indicated in Figure 10(b). The details in this case are similar to the previous case except that now the two crossings corresponding to $\{y_i, y_i^{-1}\}$ occur on the same side of the z -hook and we have the following possibility: Suppose that c does not contain any type (II) canceling pairs in z . Then, we may have a type (II) canceling pair $\{y_j, y_j^{\pm 1}\}$, for some $j > i$, such that the crossings in C_z separate crossings corresponding y_j and $\bar{\delta}_c = \emptyset$. This corresponds to the exceptional case mentioned in the statement of the lemma. In this case, the arc representing $[y_1, \dots, y_i, \dots, y_j]$ can be seen to be a *good* arc.




Figure 11. Both the endpoints of the arc can be further hooked with the z -hook

An example of an arc where this exceptional case is realized is shown in Figure 11. Here we have $i = 1$. Notice that both endpoints of the arc shown here can be hooked with the z -hook. Thus we can form $[y_1, y, y^{-1}, \dots, y_j, \dots]$, where $y_1 = z$, $y_j = z^{\pm 1}$ and the crossings corresponding to $\{y_j, y_j^{-1}\}$ occur on different endpoints of the z -hook.

b) It follows immediately from part a) and 3.12(b).

c) Let $d = [y_1, \dots, y_{j-1}]$ and suppose $y_j = y^{\pm 1}$, for some free generator. Then $[y_1, \dots, y_j] = [d, y^{\pm 1}]$.

Let δ_1 be the arc representing $[d, y^{\pm 1}]$. A moment's thought will convince us that in order for δ_1 to be bad the following must be true: i) The arc δ_1^* of Lemma 3.10 must intersect the y -hook precisely once; and ii) crossings on the y -hook, corresponding to some appearance of $y^{\pm 1}$ in d , must separate the crossings corresponding to the canceling pair $\{y_j, y_j^{-1}\}$. In particular, $y^{\pm 1}$ must have appeared in d at least once. Moreover it follows from Lemma 3.8 that, for any commutator c , in order to be able to form $[[c, y^{\pm 1}], y^{\pm 1}]$, $[c, y^{\pm 1}]$ must be represented by a good arc. Thus we may assume that d satisfies the following: at least one of the y_i 's is equal to $y^{\pm 1}$; and $y_{j-1} \neq y^{\pm 1}$.

Case 1: Suppose that the arc δ_1 representing $[d, y_j]$ doesn't run through the hook of the band B ; in particular δ_1 is embedded. Then, any canceling pair in d is admissible. From our assumption above, the only remaining possibility is when $\{y_j, y_j^{-1}\}$ corresponds to the exceptional case of part a). As already said in the proof of a), in this case δ_1 is good.

Case 2: Suppose that δ_1 runs through the hook of B . Then,

$$[d, y_j] = [e, y_r, \dots, y_{j-1}, y_j],$$

where i) $\{y_r, y_r^{-1}\}$ is the special canceling pair, and δ_1 runs through the hook at this stage and ii) $e = [y_1, \dots, y_{r-1}]$ is a simple (quasi-)commutator.

Notice that all the canceling pairs corresponding to y_k with $k \neq r$ are admissible, by definition. Let x_0 be the free generator corresponding to the band containing δ . By Lemma 3.8, we see that either $y_r = x_0^{\pm 1}$ or the arc representing $[d, y_j]$ is embedded. In the later case, it follows by Lemma 3.12 a), that δ_1 is embedded and the result follows as in Case 1. If $y_r = x_0^{\pm 1}$, then all the sub-arcs of δ_1 representing e or e^{-1} are embedded; a moment's thought will convince us

that are good arcs. Thus if $j = r$, the conclusion of the lemma follows. Suppose $j > r$. By 3.12 a) we have $y_i \neq x_0^{\pm 1}$, for all $i > r$. In particular, $y \neq x_0^{\pm 1}$. Now the conclusion follows as in Case 1. $\square$

Before we are ready to state and prove the result about good arcs promised in the beginning of 3c, we need some more notation and terminology.

Let $c = [y_1, y_2, \dots, y_{n-1}, y_n]$ and let $\mathcal{C} = \{C_1, C_2, \dots, C_{n-1}, C_n\}$ the sets of letters of Lemma 2.10. We will denote by $||\delta||$ the cardinality of the maximal subset of $\mathcal{C}$ that trivializes δ geometrically; that is δ is $||\delta||$ -nice (see Definition 3.4). We will denote by $s(c)$ the number of bad sets in $\mathcal{C}$. For a quasi-commutator $\hat{c}$, we will define $s(\hat{c}) = s(c)$ where c is the commutator underlying $\hat{c}$.

Finally, for $n \in \mathbb{N}$, let $t(n)$ be the quotient of the division of n by four, and let $q(n)$ be the quotient of division by six.

Lemma 3.15. *Suppose that S, B, γ and δ are as in the statement of Proposition 3.3, and that δ_1 is a good subarc of δ representing a simple quasi-commutator c_1 , of length $m + 1$.*

a) If δ_1 is embedded then δ_1 is an $t(m + 1)$ -nice arc.

b) If δ_1 is non embedded then δ_1 is an $q(m + 1)$ -nice arc.

Proof. a) Inductively we will show that

$$||\delta_1|| \geq m + 1 - s(c_1) \tag{1}$$

Before we go on with the proof of (1), let us show that it implies that δ_1 is $t(m)$ -nice.

For a fixed free generator y , let $w(y)$ be the number of appearances of y in c_1 and let $s^y(c_1)$ be the number of bad sets in y . By Lemma 3.14 a), with one exception, a set C_i corresponding to y can become bad only by a successive appearance of y . By Lemma 3.12, no letter can appear in c_1 more than two successive times. A simple counting will convince us that

$$\frac{w(y)}{s^y(c_1)} \geq \frac{4}{3},$$

and that the maximum number of bad sets in a word is realized when each generator involved appears exactly four times, three of which are bad. Thus we have $s(c_1) \leq m + 1 - t(m + 1)$ and by (1) we see that $||\delta_1|| \geq t(m + 1)$, as desired.

We now begin the proof of (1), by induction on m . For $m = 1$, we know that all (embedded) arcs representing a simple 2-commutator are nice and thus (1) is true. Assume that $m \geq 2$ and (inductively) that for every good arc representing a commutator of length $\leq m$, (1) is satisfied. Now suppose that

$$c_1 = [[c, z^{\pm 1}], y^{\pm 1}]$$

where c is a simple commutator of length $m - 1$, and z, y are free generators.

Let $\bar{\delta}_{1,2}$ (resp. $\bar{\theta}_{1,2,3,4}$) denote the subarcs of δ_1 representing $[c^{\pm 1}, z^{\pm 1}]^{\pm 1}$ (resp. $c^{\pm 1}$).

Case 1. The arcs $\bar{\delta}_{1,2}$ are good. By induction we have

$$||\bar{\delta}_{1,2}|| \geq m - s(\bar{c}), \quad (2)$$

where $\bar{c} = [c, z^{\pm 1}]$. Since δ_1 is good, a set of crossings that trivializes $\bar{\delta}_{1,2}$ can fail to work for δ_1 only if it becomes a bad set in c_1 . Moreover, the set of crossings corresponding to the last canceling pair $\{y^{\pm 1}, y^{\mp 1}\}$ of c_1 , also trivializes δ_1 geometrically. By 3.14a) forming c_1 from $[c, z^{\pm 1}]$ can create at most two bad sets, each involving $y^{\pm 1}$. Thus we have $s(\bar{c}) \leq s(c_1) \leq s(\bar{c}) + 2$. Combining all these with (2), we obtain

$$||\delta_1|| \geq ||\bar{\delta}_{1,2}|| + 1 - 2 \geq m + 1 - s(\bar{c}) - 2 \geq m + 1 - s(c_1),$$

which completes the induction step in this case.

Case 2. Suppose that $\bar{\delta}_{1,2}$ are not good arcs. Let us use $\bar{\theta}$ to denote any of $\bar{\theta}_{1,2,3,4}$. Suppose that $c = [c_2, x]$, and thus $c_1 = [[[c_2, x], z^{\pm 1}] y^{\pm 1}]$. By Lemma 3.14(c), $\bar{\theta}$ is a good arc and by induction

$$||\bar{\theta}|| \geq m - 1 - s(c). \quad (3)$$

First suppose that $y \neq z^{\pm 1}$. Because $\bar{\delta}_{1,2}$ are not good we can't claim that the pair $\{z^{\pm 1}, z^{\mp 1}\}$ trivializes $\bar{\delta}_{1,2}$ geometrically; however it will work for δ_1 . Moreover, the set of crossings corresponding the last canceling pair $\{y^{\pm 1}, y^{\mp 1}\}$ of c_1 , also trivializes δ_1 geometrically. Notice that the only sets of crossings that work for $\bar{\theta}$ but could fail for δ_1 are these involving $z^{\pm 1}$ or $y^{\pm 1}$ that correspond to bad pairs in δ_1 . We see that

$$s(c) \leq s(c_1) \leq s(c) + 4.$$

Combining all these with (3), we obtain

$$||\delta_1|| \geq ||\bar{\theta}|| + 2 - 4 \geq m + 1 - s(c_1),$$

which completes the induction step in this case.

Now suppose that $y = z$. In this case we can see that $s(c) \leq s(c_1) \leq s(c) + 2$ and that at least one of the two last canceling pairs c_1 will trivialize δ_1 geometrically. These together with (3) imply (1). This finishes the proof of part a) of our lemma.

b) Let x_0 denote the free generator of π corresponding to it. If c_1 doesn't involve x_0 at all, δ_1 has to be an embedded good arc and the conclusion follows from part a). So we may suppose that δ_1 involves x_0 . Now the crossings that correspond to appearances of x_0 in c_1 may fail to trivialize the arc geometrically.




Figure 12. Simple commutators occupying the entire band

See, for example the arcs in Figure 12; in both cases crossings that realize the contributions of x_0 fail to trivialize the band. As a result of this, we can only claim that

$$||\delta_1|| \geq m + 1 - (s^*(x_0) + w(x_0)), \quad (4)$$

where $w(x_0)$ denotes the number of appearances of x in c_1 and $s^*(x_0)$ is the number of the bad sets in generators different than x_0 .

The proof of (4) is similar to that of (1) in part a). Now by Lemmas 3.12 and 3.14 it follows that the word c_1 that will realize the maximum number of bad sets has the following parity:

$$[x_0, y_1, y_1, x_0, x_0, \dots, y_k, y_k, x_0, x_0, y_1, y_1, \dots, y_k, y_k],$$

where $y_1, \dots, y_k$ are distinct and $x_0 \neq y_i$. Moreover, three out of the four appearances of each y_i correspond to bad sets. Now the conclusion follows. $\square$

Remark 3.15*. From the proof of Lemma 3.15 we see that if the arc δ_1 realizes the maximum number of bad sets then $q(m+1) = k$; the number of distinct generators, besides x_0 , involved in c_1 . In fact, a careful reading of the proof of 3.15 will reveal that

$$\|\delta_1\| \geq \begin{cases} q(m+1), & \text{if } m < 6k \\ k + \left\lceil \frac{m-6k}{2} \right\rceil, & \text{if } m \geq 6k. \end{cases}$$

Thus $\|\delta_1\| \geq q_{\delta_1}$, and Proposition 3.3 follows for bands representing a simple (quasi-)commutator.

3d. Conflict sets and products of good arcs

In this paragraph we study arcs that decompose into products of good arcs. The main result of the paragraph is Lemma 3.19, in which we show that an arc $\tilde{\delta}$ that is a product of good arcs is $q_{\tilde{\delta}}$ -nice.

First, we need some more notation and terminology. Let S, B, γ and δ be as in the statement of Proposition 3.3, and let $\tilde{\delta}$ be a subarc of γ representing a word W_1 in $\pi^{(m+1)}$. Suppose that $W_1 = \hat{c}_1 \dots \hat{c}_s$, is a product of quasi-commutators represented by arcs $\{\delta_1, \dots, \delta_s\}$, respectively. Also, for $k = 1, \dots, s$, let $\mathcal{C}_k = \{C_1^k, \dots, C_{(m+1)}^k\}$ be the sets of crossings of Lemma 2.10 for δ_i . Let $C \in 2^{\mathcal{C}_k}$; by assumption the set of letters in C trivialize W_1 algebraically. For a proper subset $\mathcal{D} \subset \{\delta_1, \dots, \delta_s\}$ we will use $C \cap \mathcal{D}$ (resp. $C \cap \bar{\mathcal{D}}$) to denote the crossings in C that lie on arcs in $\mathcal{D}$ (resp. in $\bar{\mathcal{D}}$). Here, $\bar{\mathcal{D}}$ denotes the complement of $\mathcal{D}$ in the set $\{\delta_1, \dots, \delta_s\}$.

For every free generator, say y , we may have crossings on the y -hook, realizing letters in the word W_1 , that trivialize geometrically some of the subarcs δ_i but fail to trivialize $\tilde{\delta}$. To illustrate how this can happen, consider the arcs δ_1 and δ_2 . Let C^1 and C^2 be sets of crossings, on the y -hook, along δ_1 and δ_2 , respectively. Suppose that C^i trivializes δ_i geometrically (i.e. it is a good set of crossings). Suppose, moreover, that there are crossings on δ_2 corresponding to a canceling pair $\{y^{\pm 1}, y^{\mp 1}\}$ that doesn't belong in C^2 , and such that they are separated by crossings in C^1 . Then $C^1 \cup C^2$ may not trivialize $\delta_1 \cup \delta_2$. With the situation described above in mind, we give the following definition.

Definition 3.16. *A set C of crossings on $\tilde{\delta}$ is called a conflict set iff i) the letters in C trivialize W_1 algebraically; ii) switching the crossings in C doesn't trivialize $\tilde{\delta}$ geometrically; and iii) there exists a proper subset $\mathcal{D}_C \subset \{\delta_1, \dots, \delta_s\}$ such that $C \cap \mathcal{D}_C$ trivializes geometrically the union of arcs in $\mathcal{D}_C$ and $C \cap \bar{\mathcal{D}}_C$ trivializes geometrically the union of arcs in $\bar{\mathcal{D}}_C$.*

Before we are ready to treat compositions of good arcs we need two auxiliary lemmas that describe the situations that create conflict sets.

Lemma 3.17. *For $k = 1, \dots, s$, let $\hat{c}_k$ be simple (quasi-)commutator represented by an arc δ_k , and let $\{C_1^k, \dots, C_{(m+1)}^k\}$ be sets of crossings as above. Moreover let $[y_1^k, y_2^k, y_3^k, \dots, y_{m+1}^k]$ be the underlying commutator of $\hat{c}_k$. Suppose that for some $i = 1, \dots, m+1$, $y_i^k = y^{\pm 1}$ for some free generator y , and that $C_i = \cup_{k=1}^s C_i^k$, is a conflict set. Let $\mathcal{D}_{C_i}$ be as in Definition 3.16. Then, there exist arcs $\delta_t \in \mathcal{D}_{C_i}$ and $\delta_r \in \bar{\mathcal{D}}_{C_i}$ such that the crossings on the y -hook corresponding to canceling pairs in y_j^r on δ_r , are separating by crossings corresponding to y_i^t on δ_t . Here $j \neq i$.*

Proof. It follows from Definitions 2.3, 3.4 and 3.16. $\square$

With the notation as in Lemma 3.17, the set C_j^r will be called a *conflict partner* of C_i^t .

Lemma 3.18. *Let $W_{1,2} = [y_1^{1,2}, y_2^{1,2}, y_3^{1,2}, \dots, y_{m+1}^{1,2}]$ be the underlying commutators of quasi-commutators represented by arcs $\delta_{1,2}$. Let C_i^1 and C_j^2 be sets of letters in W_1 and W_2 respectively, corresponding to the same free generator y . Suppose that C_j^2 is a conflict partner of C_i^1 . Then, with at most one exception, (1) either $j = i + 1$ (resp. $j = i - 1$) and $y_k^1 = y_k^2$ for $k < i$ (resp. for $k < i - 1$); or*

(2) the sets of free generators appearing in $\{y_1^1, \dots, y_i^1\}$ (resp. $\{y_1^1, \dots, y_j^1\}$) and $\{y_{i+1}^2, \dots, y_{j-1}^2\}$ (resp. $\{y_{j+1}^2, \dots, y_{i-1}^2\}$), are disjoint.

Proof. By 3.17, there must be crossings on the y -hook corresponding to canceling pairs in C_j^2 , that are separated by crossings in C_i^1 . Let C_1 denote the canceling pair corresponding to y_i^1 in $[y_1^1, y_2^1, \dots, y_i^1]$. and let C_2 denote the canceling pair corresponding to y_i^2 (resp. y_j^2) in $[y_1^2, y_2^2, \dots, y_i^2]$ (resp. $[y_1^2, y_2^2, \dots, y_j^2]$) if $j > i$ (resp. $j < i$). By Lemma 3.12(a), $C_{1,2}$ are of type (I) or (II). Let $D_{1,2}$ be the finite disc corresponding to the canceling pair $C_{1,2}$, in $W_{1,2}$, respectively. Up to symmetry there are three cases to consider: i) Both $C_{1,2}$ are of type (I); ii) both $C_{1,2}$ are of type (II); and iii) one of them is of type (I) and the other of type (II). In each case the result will follow using Digression 3.9 to study the components of $D_{1,2} \cap \bar{\delta}_{1,2}$, where $\bar{\delta}_{1,2}$ denotes the complement in $\delta_{1,2}$ of the parallel arcs corresponding to $C_{1,2}$, respectively. The exceptional case will occur when the canceling pair C_1 is of type (I) and crossings in it are separated by a type (II) canceling pair on δ_2 . The details are similar to these in the proof part a) of Lemma 3.14 $\square$

To continue, recall the quantity $q_{\tilde{\delta}}$ defined before the statement of Proposition 3.3.

Lemma 3.19. (Products of good arcs) *Let S , B , γ and δ be as in the statement of Proposition 3.3, and let $W = c_1 \dots c_r$ be a word expressing δ^+ as a product of simple quasi-commutators. Suppose that $\tilde{\delta}$ is a subarc of δ , representing a subword of simple quasi-commutators $W_1 = \hat{c}_1 \dots \hat{c}_s$, each of which is represented by a good arc. Then $\tilde{\delta}$ is $q_{\tilde{\delta}}$ -nice. In particular if W is a product of simple quasi-commutators represented by good arcs, Proposition 3.3 is true for δ .*

Proof. If $s = 1$ the conclusion follows from Lemma 3.15 and Remark 3.15*. Assume that $s > 1$. Let $\delta_1, \dots, \delta_s$ be arcs representing $\hat{c}_1 \dots \hat{c}_s$, respectively.

In general, we may have conflict sets of crossings between the δ_j 's. Since conflict sets occur between commutators that have common letters, we must partition the set $\{\hat{c}_1, \dots, \hat{c}_s\}$ into groups involving disjoint sets of generators and work with each group individually. The maximum number of conflicts will occur when all the $\hat{c}_i$'s belong in the same group. Since conflict sets are in one to one correspondence with proper subsets of $\{\delta_1, \dots, \delta_s\}$, the maximum number of conflict sets, for a

fixed generator y , is $2^s - 2$.

Let x_0 be the generator corresponding to the hook of B . From the proof of Lemma 3.15, Remark 3.15* and by Lemma 3.18 we can see that a word W , in which there are k distinct generators besides $x_0^{\pm 1}$, will realize the maximum number of bad sets of crossings on the individual δ_i 's and the maximum number of conflict sets, if the following are true:

- i) The length $m + 1$ is equal to $6k + r + 2k(2^s - 2)$, where $r > 2$;
- ii) each of the arcs δ_i realizes the maximum number of bad sets and the maximum number of appearances of $x_0^{\pm 1}$ (i.e. $5k + \frac{r}{2}$) and there are $k(2^s - 2)$ conflict sets between the δ_i 's. Moreover, each pair of conflict partners in W correspond either to the exceptional case or in case (1) of Lemma 3.18.

We claim, however, that there will be $k + \left\lceil \frac{r}{2} \right\rceil + k(s - 2)$ sets of crossings that trivialize $\tilde{\delta}$ geometrically. From these $k + \left\lceil \frac{r}{2} \right\rceil$ come from good sets on the δ_i 's. The rest $ks - 2k$ are obtained as follows: For a fixed $y \neq x_0^{\pm 1}$, the crossings in the conflict sets involving $y^{\pm 1}$ and in their conflict partners can be partitioned into $s - 2$ disjoint sets that satisfy the definition of $(s - 3)$ -triviality. To see that, create an $s \times (2^s - 2)$ matrix, say A , such that the (i, j) entry in A is the j -th appearance of y in $\hat{c}_i$. The columns of A are in one to one correspondence with the conflict sets $\{C_i\}$, in y . By 3.18, and 3.12 there are at most $2s$ "exceptional" conflict partners shared among the C_i 's. Other than that, the conflict partners of a column C_i will lie in exactly one of the adjacent columns. For $s \geq 4$ we have $2^s - 2 \geq 4s$ and thus A has at least $2s$ columns that can only conflict with an adjacent column; these will give $s > s - 2$ sets as claimed above. For $s = 2, 3$ the conclusion is trivial.

Now from i) we see that $ks - 2k > \log_2\left(\frac{m + 1 - 4k - r}{4}\right)$. Thus,

$$\frac{r}{2} + k(s - 2) > \log_2\left(\frac{m + 1 - 4k}{4}\right) > \log_2\left(\frac{m + 1 - 6k}{6}\right),$$

and the claim in the statement of the lemma follows. $\square$

3e. The reduction to nice arcs

Let S , B , γ and δ be as in the statement of Proposition 3.3. Our goal in this paragraph is to finish the proof of 3.3. We begin with the following lemma, which relates the q_μ -niceness of a subarc $\mu \subset \tilde{\delta} \subset \delta$ to the $q_{\tilde{\delta}}$ -niceness of $\tilde{\delta}$.

Lemma 3.20. (Products of nice arcs) *Let S , B , γ and δ be as in the statement of Proposition 3.3. Let $\tilde{\delta}$ be a subarc of δ , representing a subword $W_1 = \hat{c}_1 \dots \hat{c}_s$, where $\hat{c}_i$ is a product of simple quasi-commutators represented by an arc θ_i . Suppose that θ_i is q_{θ_i} -nice, for $i = 1, \dots, s$. Then $\tilde{\delta}$ is $q_{\tilde{\delta}}$ -nice.*

Proof. Once again we can have sets of crossings on $\tilde{\delta}$ that trivialize W_1 , and trivialize a subset of $\{\theta_1, \dots, \theta_s\}$ geometrically but fail to trivialize $\tilde{\delta}$. For $i = 1, \dots, s$, let m_i denote the number of simple quasi-commutators in $\hat{c}_i$, and let $\mathcal{D}_i$ denote the set of subarcs of $\tilde{\delta}$ representing them. We notice that the maximum number of conflict sets that we can have in W_1 , is $k[2^{(m_1 + \dots + m_s)} - 2]$ where k is the number of distinct generators, different than x_0 , appearing in W . Now we may proceed as in the proof of Lemma 3.19. $\square$

To continue recall the notion of a quasi-nice arc (Definition 3.4). Our last lemma in this section shows that the notions of quasi-nice and $q_{\tilde{\delta}}$ -nice are equivalent.

Lemma 3.21. *A quasi-nice subarc $\tilde{\delta} \subset \delta$ that represents a product $W = c_1 \dots c_r$ of quasi-commutators, is $q_{\tilde{\delta}}$ -nice.*

Proof. Let $\delta_1, \dots, \delta_r$ be the arc representing $c_1, \dots, c_r$, respectively. Let $\mathcal{D}_g$ (resp. $\mathcal{D}_b$) denote the set of all good (resp. not good) arcs in $\{\delta_1, \dots, \delta_r\}$. Also let n_g (resp. n_b) denote the cardinality of $\mathcal{D}_g$ (resp. $\mathcal{D}_b$). If $n_b = 0$, the conclusion follows from Lemma 3.19. Otherwise let $\mu \in \mathcal{D}_b$, be the first of the δ_i 's not represented by a good arc. Suppose it represents $c_\mu = [c^{\pm 1}, y^{\mp 1}]$, where $c^{\pm 1}$ is a simple quasi-commutator of length m , and y a free generator. Let μ^1, μ^2 be the subarcs of μ representing $c^{\pm 1}$. Since μ is not good, y must have appeared in c ; thus the numbers of distinct generators in the words representing μ and μ^1, μ^2 are the same. We see that $q_{\mu^1} = q_{\mu^2} = q_\mu$. By 3.14, μ^1, μ^2 are good arcs and by 3.15 they are q_μ -nice. Let $\bar{\mu} = \tilde{\delta} \setminus \mu$, and let $\bar{\mu}_{1,2}$ denote the two components of $\bar{\mu}$. By induction and 3.20, $\bar{\mu}_i$ is $q_{\bar{\mu}_i}$ -nice. Since μ is not good, one of its endpoints lies inside the y -hook and the other outside. Moreover the arc μ^* of Lemma 3.10, separates crossings corresponding to canceling pairs on the y -hook. Now a moment's thought will convince us that at least one of $\bar{\mu}_{1,2}$ must have crossings on the y -hook. A set of crossings that trivializes geometrically $\theta_1 = \mu^1 \cup \mu^2$ and $\theta_2 = \bar{\mu}_1 \cup \bar{\mu}_2$ will fail to trivialize $\tilde{\delta}$ only if there are conflict sets between θ_1 and

θ_2 . A counting argument shows that the maximum number of conflict sets that can be on $\tilde{\delta}$ is $k(2^r - 2)$, where k is the number of distinct generators, different than x_0 , in W . Now the conclusion follows as in the proof of 3.20. $\square$

Proof of Proposition 3.3. It follows immediately from 3.21 and the fact that the arc δ in the statement of 3.3 is quasi-nice; see discussion at the end of 3b. $\square$

Remark 3.22. a) Theorem 3.2 is not true if we don't impose any restrictions on the surface S of Definition 3.1. For example let K be a positive knot set $\pi_K = \pi_1(S^3 \setminus K)$ and let D_K denote the untwisted Whitehead double of K . Let S be the standard genus one Seifert surface for $D(K)$. Since $\pi_K^{(n)} = \pi_K^{(2)}$ for any $n \geq 2$, we see that S has a half basis realized by a curve that if pushed in the complement of S lies in $\pi_K^{(n)}$, for all $n \geq 2$. On the other hand, D_K doesn't have all its Vassiliev invariants trivial since it has non-trivial 2-variable Jones polynomial (see for example [Ru]).

b) The methods of this section were applied in [K-L] to obtain relations between finite type and Milnor invariants of links that are realized as plat-closures of pure braids.

4. VASSILIEV INVARIANTS AS OBSTRUCTIONS TO n -SLICENESS

Let S be a *regular* Seifert surface. In this section we show that the Vassiliev invariants of the knot $K = \partial S$ are null-concordance obstructions of links that can be derived from *regular* spines of S .

Let B^4 denote the 4-ball, and let $L = L_1 \cup \dots \cup L_m$ be an m -component link in $S^3 = \partial B^4$. We begin by recalling, from [Co] or [O], the definition of n -slice (or n -null-cobordant).

Definition 4.1. ([Co], [O]) *The link L is called n -slice if there exist m disjoint connected surfaces $V = V_1 \cup \dots \cup V_m \subset B^4$, with $\partial V_i = L_i$ and such that the following is true: For some trivialization of a tubular neighborhood of V in B^4 , which restricts to the standard trivialization of a tubular neighborhood of L in $S^3 = \partial B^4$, the composition*

$$\pi_1(V_i) \longrightarrow \pi_V \longrightarrow \pi_V / \pi_V^{(n)}$$

is trivial, for all i . Here, $\pi_V = \pi_1(B^4 \setminus V)$.

As was shown in [Co] and [O], n -slice is the geometric notion which is equivalent to the vanishing of Milnor's invariants. More precisely we have:

Theorem 4.2. ([Co], [O]) *A link $L = L_1 \cup \dots \cup L_m$ is n -slice for all $n \in \mathbb{N}$ if and only if all its Milnor invariants vanish. That is, the homomorphism $F \longrightarrow \pi_L$, given by any choice of meridians of L , induces isomorphisms*

$$F/F^{(k)} \longrightarrow \pi_L/\pi_L^{(k)}$$

for all $k \leq n + 1$ and all $n \in \mathbb{N}$. Here F is the free group of rank m , and $\pi_L = \pi_1(S^3 \setminus L)$.

To continue, let S be a regular Seifert surface of a knot K , and let Σ be a regular spine of S , consisting of curves $\gamma_1, \beta_1, \dots, \gamma_g, \beta_g$. Let

$$\bar{\epsilon} := (\epsilon_1, \dots, \epsilon_{2g})$$

where each ϵ_i is equal to $+$ or $-$. Finally, let

$$L^{\bar{\epsilon}} =: \gamma_1^{\epsilon_1} \cup \beta_1^{\epsilon_2} \cup \dots \cup \beta_g^{\epsilon_{2g}}$$

Clearly, $L^{\bar{\epsilon}}$ is a link with $2g$ -components. Let $\pi = \pi_1(S^3 \setminus S)$ and let $\pi_{L^{\bar{\epsilon}}} = \pi_1(S^3 \setminus L^{\bar{\epsilon}})$. By assumption, π is the free group of rank $2g$. We will say that the spine Σ is *admissible* if the following is true: For some $\bar{\epsilon}$ as above, the longitudes of $L^{\bar{\epsilon}}$ lie in $\pi_{L^{\bar{\epsilon}}}^{(n)}$ (for some $n \in \mathbb{N}$) if and only if $\gamma_i^{\epsilon_i}, \beta_i^{\epsilon_{i+1}} \in \pi^{(n)}$. The link $L^{\bar{\epsilon}}$ will be called an *admissible spine link* for S . The theorem below shows the Vassiliev invariants of the boundary knot are obstructions of null-cobordance for the links $L^{\bar{\epsilon}}$.

Theorem 4.3. *Let S be a regular Seifert surface of a knot K and suppose that $L^{\bar{\epsilon}}$ is an admissible spine link for S . If the Milnor invariants of length $\leq n + 1$ vanish for $L^{\bar{\epsilon}}$ then all the Vassiliev invariants of orders $\leq l(n)$ vanish for K .*

Proof. It follows from the fact that if the Milnor invariant of length $\leq n + 1$ vanish for $L^{\bar{\epsilon}}$ then its longitudes lie in $\pi_{L^{\bar{\epsilon}}}^{(n+1)}$ ([M2]), our discussion above, and Theorem 3.2. $\square$

It was conjectured (see [Co]) that a link L is n -slice for some $n \in \mathbb{N}$ if and only if Milnor's invariants with length less or equal to $2n$ vanish. For the proof of one direction of this conjecture (namely, that n -slice implies the vanishing of invariants of length $\leq 2n$) see [L2]. The proof of the other direction was given by Igusa and Orr ([I-O]). Thus we obtain.

Corollary 4.4. *If L^ε is n -slice then the Vassiliev invariants of orders $\leq l(2n - 1)$ vanish for K .*

Remark 4.5. It is known (see [N-S] for relevant discussion) that for every $n \in \mathbb{N}$, and every non-slice knot K one can find slice (null-concordant) knots whose Vassiliev invariants of orders $\leq n$ are equal to these of K . It is interesting to compare Theorem 4.3 (or Corollary 4.4) with this fact.

5. MORE TYPES OF n -TRIVIAL KNOTS

Using the calculation of the Alexander polynomial via Seifert matrices one can see that the Alexander polynomial of an n -hyperbolic knot is trivial. On the other hand there are n -trivial knots with non-trivial Alexander polynomial. Our purpose in this section is to generalize the notion of n -hyperbolic so that we can include knots with non-trivial Alexander polynomial. The generalized notion is that of n -unknotted (see Definition 5.5). Roughly speaking, a knot is called n -unknotted if it bounds a Seifert surface whose complement looks, modulo certain terms of its fundamental group, “simple” (n -parabolic). We show that the existence of such a surface for a knot implies the vanishing of its Vassiliev invariants of orders $\leq n$. See Theorem 5.6. To complement the notions of n -hyperbolic and n -parabolic we also distinguish another special class of n -unknotted; namely that of n -elliptic. We will have more to say about each of these classes of knots and their relation with each other in §6.

5a. Definitions

We begin by introducing some notation and terminology needed to continue. Let K be a knot and let S be a genus g regular Seifert surface of K , in disc-band form. For a band A of S let γ_A and x_A denote the core and the free generator of $\pi = \pi_1(S^3 \setminus S)$ corresponding to A , respectively. Also, for a set of bands $\mathcal{A}$ we will

denote by $\mathcal{G}_{\mathcal{A}}$ (resp. $\mathcal{D}_{\mathcal{A}}$) the normal subgroup of π generated by $\{x_A \mid A \in \mathcal{A}\}$ (resp. $\{x_A \mid A \in \mathcal{CA}\}$). Here $\mathcal{CA}$ denotes the complement of $\mathcal{A}$.

We will say that two bands A, A' are *geometrically related* if there exists a sequence of bands

$$A = A_0 \rightarrow A_1 \rightarrow \dots \rightarrow A_{r-1} \rightarrow A_r = A',$$

such that, for $i = 1, \dots, r$, $\gamma_{A_i}^+$ or $\gamma_{A_i}^-$ is geometrically linked to at least one of $\gamma_{A_0}, \dots, \gamma_{A_{i-1}}$. Otherwise A and A' will be called *geometrically unrelated*. Moreover, we will say that two sets of bands $\mathcal{A}$ and $\mathcal{B}$ are *geometrically unrelated* if any two $A \in \mathcal{A}$ and $B \in \mathcal{B}$ are geometrically unrelated.

For a band A whose core γ_A represents an element in $\pi^{(m+1)}$ let q_{γ_A} be the quantity defined in the beginning of §3.

Definition 5.1. Let $n \in \mathbb{N}$, with $n > 1$. A regular Seifert surface S will be called *n-elliptic*, iff there exist two half bases $\mathcal{A}$ and $\mathcal{B}$, represented by circles in a regular spine Σ , such that the following is true: The sets $\mathcal{A}$ and $\mathcal{B}$ are geometrically unrelated and for every $A \in \mathcal{A}$ there exist $B \in \mathcal{B}$ and $m_A, m_B \in \mathbb{N}$ with $q_{\gamma_A} + q_{\gamma_B} = n + 1$ and such that we have

$$[\gamma_A^\epsilon] \in \mathcal{G}_{\mathcal{A}}^{(m_A+1)}$$

and

$$[\gamma_B^{-\epsilon}] \in \mathcal{G}_{\mathcal{B}}^{(m_B+1)}$$

The boundary of such a surface will be called an *n-elliptic knot*.

Definition 5.2. Let $n \in \mathbb{N}$, with $n > 1$. A regular Seifert surface S will be called *n-parabolic*, iff there exist two half bases $\mathcal{A}$ and $\mathcal{B}$, represented by circles in a regular spine Σ , such that the following is true:

- a) The sets $\mathcal{A}$ and $\mathcal{B}$ are geometrically unrelated. Moreover, any two subsets of $\mathcal{A}$ are geometrically unrelated.
- b) Let $s = s(K, S) \geq 1$ be the largest integer such that K can be unknotted in $2^s - 1$ distinct ways by untwisting along the bands in $\mathcal{A}$. If $n > s$, for every $B \in \mathcal{B}$ we have that

$$[\gamma_B^\epsilon] \in \mathcal{G}_{\mathcal{B}}^{(m+1)}$$

where $m \in \mathbb{N}$, such that with $q_{\gamma_B} + s = n + 1$.

The boundary of such a surface will be called an n -parabolic knot. The numbers s and $t = n + 1 - s$ will be called the simplicity and triviality of K , respectively.

Remark 5.3. Our terminology for regular surfaces, in Definitions 3.1, 5.1 and 5.2 has been motivated by the forms of their Seifert matrices. To explain this, let S be a regular surface and let M be the matrix $V + V^T$, where V is a Seifert matrix of S . We see that if S is n -elliptic, then there exists a regular basis of $H_1(S)$, with respect to which M is of the form

$$\begin{bmatrix} J & 0 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & J \end{bmatrix}$$

where

$$J = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Thus M is an elliptic matrix.

If S is n -hyperbolic then M is congruent to a matrix of the form

$$\begin{bmatrix} 0 & P \\ P & R \end{bmatrix}$$

for some $g \times g$ matrices P, R . Finally, if S is n -parabolic then M is congruent to a matrix of the form

$$\begin{bmatrix} D_g & D \\ D & T \end{bmatrix}$$

where D_g and D are diagonal $g \times g$ matrices and $T = 0$ if $n \gg 0$.

We have:

Theorem 5.4. *An n -elliptic or n -parabolic knot is n -trivial.*

Proof. In both cases the proof will be by induction on the genus of the Seifert surface, with the desired properties.

a) First we assume that K is n -elliptic.

If the genus of the surface in Definition 5.1 is zero, the conclusion follows trivially. Otherwise, let us concentrate on a pair of dual bands (A, B) . We can assume that $[\gamma_A^\epsilon] \neq 1$ and $[\gamma_B^{-\epsilon}] \neq 1$.

Since by assumption the sets $\mathcal{A}$ and $\mathcal{B}$ are *geometrically unlinked* we may modify the projection of S so that Proposition 3.3 applies to both bands, simultaneously. To see this, first we find a projection with respect to which the

requirements of Lemma 2.12 are satisfied for A . Let l be the horizontal line associated to this projection. Then, we start sliding the overcrossings of B that are below l (see Remark 2.5) till B is put in good position. At the end of the procedure, our projection will have the following property: Every crossing between A (resp. B) and any other band that occurs below l will be an undercrossing (resp. overcrossing).

These properties together with Proposition 3.3 guarantee the following: We have $n + 1$ sets of crossings ($C_1, \dots, C_{q(m_A)}$ from A and $C_{q(m_A)+1}, \dots, C_{n+1}$ from B) that enable us to write K , in $\mathcal{V}_n$, as a linear combination of $2^{n+1} - 1$ knots $\{K_C\}$, with the following property: each K_C is an n -elliptic knot that bounds an n -elliptic Seifert surface, of genus less than g . Thus the induction step applies.

b) Now we suppose that K is n -parabolic.

Let S be an n -parabolic Seifert surface for K , and let g , s and t be the genus, the simplicity and the triviality of S , respectively. By Definition 5.2 the twists along the bands in $\mathcal{A}$ provide us with s sets of crossings satisfying the definition of $s - 1$ -trivial. These together with the $t = n + 1 - s$ sets of crossings obtained by Proposition 3.3 for some $B \in \mathcal{B}$ will allow us to write K , in $\mathcal{V}_n$, as a linear combination of $2^{n+1} - 1$ knots $\{K_C\}$, with the following property: each K_C is either the trivial knot, or an n -parabolic knot that bounds an n -parabolic surface, of genus less than g . $\square$

5b. Mixed type knots

Let the notation be as in the beginning of §5a and let $\beta \in \mathcal{G}_{\mathcal{A}}^{(m+1)}$, for some set of bands $\mathcal{A}$ and $m \in \mathbb{N}$. Write β as a product, $W_1 \dots W_s$, of commutators in $\mathcal{G}_{\mathcal{A}}^{(m+1)}$ and partition the set $\{W_1, \dots, W_s\}$ into disjoint sets, say $\mathcal{W}_1, \dots, \mathcal{W}_t$ such that: i) $k_1 + \dots + k_t = l$, where k_j is the number of distinct generators involved in $\mathcal{W}_j$ and ii) for $a \neq b$, the sets of generators appearing $\mathcal{W}_a$ and $\mathcal{W}_b$ are disjoint. Let

$$k = \min\{k_1, \dots, k_t\}.$$

We define

$$q_\beta = \begin{cases} q(n+1), & \text{if } n < 6k \\ k + \left\lceil \log_2\left(\frac{n+1-6k}{6}\right) \right\rceil, & \text{if } n \geq 6k. \end{cases}$$

Before we can state the main result in this section, which is a generalization of Theorems 3.2 and 5.4, we need the following:

Definition 5.5. Let $n \in \mathbb{N}$, with $n > 1$. We will say that the knot K is n -unknotted iff it has a regular Seifert surface S , that contains two half bases $\mathcal{A}$ and $\mathcal{B}$ represented by circles in a regular spine Σ , such that the following are true: For every $A_i \in \mathcal{A}$ there exists a $B_i \in \mathcal{B}$ so that

$$[\gamma_{A_i}^\epsilon] = (x_{A_i})^{l_{A_i}} \chi_{A_i} \mu_{A_i}$$

and

$$[\gamma_{B_i}^{-\epsilon}] = \zeta_{B_i} \chi_{B_i}$$

where $\mu_{A_i} \in \pi$, $l_{A_i} \in \mathbb{Z}$, and $\zeta_{B_i} \in \mathcal{G}_{\mathcal{B}}$. Moreover, we have the following:

- a) There exist integers $m_1, \dots, m_g$, such that $\mu_{A_i} \in \pi_{\mathcal{A}_{i-1}}^{(m_i+1)}$, and with $q_{\mu_{A_i}} = n+1$ for $i = 1, \dots, g$. Here $\pi_{\mathcal{A}_{i-1}}$ and $\mathcal{A}_i$ are as in Definition 3.1.
- b) There exist $m_{A_i}, m_{B_i} \in \mathbb{N}$ such that

$$\chi_{A_i} \in \mathcal{G}_{\mathcal{A}}^{(m_{A_i}+1)}$$

and

$$\chi_{B_i} \in \mathcal{G}_{\mathcal{B}}^{(m_{B_i}+1)},$$

with $q_{\chi_{A_i}} + q_{\chi_{B_i}} = n+1$.

- c) For every A_i, B_i as in a), either $\chi_{A_i} = \chi_{B_i} = 1$ or $\chi_{A_i} \neq 1$ and $\chi_{B_i} \neq 1$ and either $\zeta_{B_i} = (x_{A_i})^{l_{A_i}} = 1$ or $(x_{A_i})^{l_{A_i}} \neq 1$ and $\zeta_{B_i} \neq 1$. Furthermore, $\chi_{A_i} \neq 1$ and $\chi_{B_i} \neq 1$ if and only if $\zeta_{B_i} = \mu_{A_i} = (x_{A_i})^{l_{A_i}} = 1$.
- d) All the ζ_{B_i} 's lie in $\mathcal{G}_{\mathcal{B}}^{(m+1)}$, for some $m \in \mathbb{N}$ with the following property: Let S_C be the Seifert surface obtained by modifying S along a set of band crossings C , that trivializes all of the $\mu_{A_i}, \mu_{B_i}, \chi_{A_i}$ and χ_{B_i} . Let s be the simplicity of $\partial(S_C)$. Then, $q_{\zeta_{B_i}} + s = n+1$.

Theorem 5.6. Assume that K , is n -unknotted, for some $n \in \mathbb{N}$. Then K is n -trivial.

Proof. Assume that K and S are as in Definition 5.5 and let g be the genus of S . Suppose that S is in disc-band form and let $A_i \in \mathcal{A}$ and $B_i \in \mathcal{B}$ be a pair of bands of S whose cores correspond to γ_{A_i} and γ_{B_i} , respectively.

We will say that the pair (A_i, B_i) is an *essential pair* if we have $\chi_{A_i} \neq 1$ and $\chi_{B_i} \neq 1$.

Let a be the number of A_i 's with $\mu_{A_i} \neq 1$ and let e be the number of *essential* pairs of bands. Finally, let b be the number of the non-trivial ζ_{B_i} 's. We define the following complexity function for (K, S) :

$$d = d(K, S) = (g, b, e, a).$$

We order the complexities lexicographically, and proceed by induction on d . For $d = (0, , ,)$ the conclusion is trivially true, since K is the unknot.

Observe that if $b = 0$ then, by Definition 5.5, either K is the unknot or it can be reduced to an n -elliptic of genus less than g ; in both cases the conclusion of the theorem follows. Suppose, now, that $b > 0$. Notice that if $e > 0$ then there is a pair of bands A_i, B_i such that $[\gamma_{A_i}^\epsilon] = \chi_{A_i}$ and $[\gamma_{B_i}^{-\epsilon}] = \chi_{B_i}$. Then we can argue as in Theorem 5.4 a), to show that K is equivalent in $\mathcal{V}_n$ to a summation of n -unknotted knots with strictly less complexity. Thus we may assume that $e = 0$. Part d) of Definition 5.5 and Proposition 3.3 guarantee the existence of $n + 1$ sets of crossings that can be used to show that K is equal, in $\mathcal{V}_n$, to an alternating summation of knots K_C such that, each K_C bounds n -unknotted surface that either has genus $< g$ or it contains a band A_i such that $[\gamma_{A_i}^\epsilon] = \mu_{A_i} \in \pi^{(m_i+1)}$, with $q_{\mu_{A_i}} = n + 1$. In both cases, it follows by Proposition 3.3 that K is equal, in $\mathcal{V}_n$, to an alternating summation of knots K_C such that, each K_C is an n -unknotted with $d(K_C) < d(K)$. $\square$

6. DISCUSSION AND QUESTIONS

In §3 and §5 we studied various classes of n -trivial knots. The following Lemma, whose proof follows directly from Definition 3.1, 5.1, 5.2 and 5.5, and the discussion following clarifies the relation between these three types.

- Lemma 6.1.** a) *Let K be a $2n$ -elliptic knot. Then K is n -hyperbolic.*
b) *Let K be an n -parabolic knot with simplicity s , and assume that $n > s + 1$. Then K is $(n - s - 1)$ -hyperbolic.*
c) *Let K be a $2n$ -unknotted knot and let s be the quantity defined in definition 5.5. In $2n > s + 1$ then K is $(2n - s - 1)$ -unknotted.*

Let $\Delta_K(t)$ denote the Alexander polynomial of a knot K . Using the representation of Alexander polynomial via Seifert matrices (see, for example, [Ro]) one can see that $\Delta_K(t) = 1$ for any n -hyperbolic knot K .




Figure 13. A 2-parabolic knot

On the other hand in Figure 13 we show a 2-parabolic knot, with simplicity $s = 2$, that has nontrivial Alexander polynomial. Moreover we have:

Proposition 6.2. *For every $n \in \mathbb{N}$, there is an n -trivial knot K , whose Alexander polynomial is non trivial.*

Proof. For a knot K , let Δ_K denote its Alexander polynomial. By the Morton-Melvin expansion ([BN-G], [M-M]) for the Alexander polynomial we have

$$\frac{p(h)}{\Delta_K(e^h)} = \sum_n v_n(K) h^n$$

where h is a variable and $p(h) = \frac{e^{\frac{h}{2}} - e^{-\frac{h}{2}}}{h}$.

Moreover, each $v_n(K)$ is a *canonical* invariant of orders n , determined by the colored Jones polynomial of K . Now the result follows from the fact that given a *canonical* (determined by its weight system) Vassiliev invariant v of order $n + 1$, there is an n -trivial knot K , such that $v(K) \neq 0$ (see for example [N-S]). $\square$

Thus the Alexander polynomial of an n -trivial knot obstructs to the knots being m -hyperbolic for any $m \in \mathbb{N}$. Now let us start with a knot K such that $\Delta_K(t) = 1$. This implies that K is algebraically slice; that is for any Seifert surface S we may find a basis of $H_1(S)$ with respect to which the Seifert matrix of S is of the form

$$V = \begin{bmatrix} 0 & A \\ B & C \end{bmatrix}$$

for some $g \times g$ matrices A , B and C . By a further change of basis of $H_1(S)$, and using the condition $\Delta_K(t) = 1$, we may change V to one of the forms that the linking matrix of a 0-hyperbolic has. Thus, we see that K is “algebraically” 0-hyperbolic. This suggests that the Alexander polynomial might be the only obstruction to m -hyperbolicity of n -trivial knots. We conjecture:

Conjecture 6.3. *A knot K is n -trivial for all $n \in \mathbb{N}$ if and only if it is n -hyperbolic for all $n \in \mathbb{N}$.*

As already mentioned, the Alexander polynomial of a knot is determined by linking numbers of “spine links” of Seifert surfaces. In particular, if “enough” of these linking numbers vanish then all the derivatives of the Alexander polynomial vanish. Our Theorem 3.2 is a generalization of this fact. It asserts that if “enough” of the higher order linking numbers (certain values Fox’s higher order free derivatives) of spines vanish then all the derivatives of all the Jones type polynomials, up to certain order, vanish. This in turn suggests that one might be able to express certain Vassiliev invariants of a knot $K = \partial S$, in terms of, appropriate values of, Fox derivatives of push offs of regular spines. For example, let S be a regular surface of K and let Σ be a regular spine of S , consisting of curves $\gamma_1, \beta_1, \dots, \gamma_g, \beta_g$. Also, let $\gamma_1^{\epsilon_1}, \beta_1^{\epsilon_2}, \dots, \beta_g^{\epsilon_{2g}}$ be as in §4 and let $W_1^{\epsilon_1}, W_1^{\epsilon_2}, \dots, W_{2g}^{\epsilon_{2g}}$ be words representing them in the free group $\pi_1(S^3 \setminus S)$. One can ask the following: Do the coefficients of the Magnus expansions of the $W_i^{\epsilon_j}$ ’s provide obstructions to the vanishing of all Vassiliev invariants of K ? This question is very important for Conjecture 6.3; in fact one needs to study “modified” finite type invariants of triples (K, S, Σ) , in terms Fox derivatives of push offs of the spine Σ . We will address this question in a subsequent preprint.

It would be interesting to know to what extent our notion of n -unknotted captures the vanishing of Vassiliev’s invariants of bounded orders. We ask the following:

Question 6.4. *Is it true that all the Vassiliev invariants of orders $\leq n$ of a knot K vanish, if and only if it is k -unknotted for some $k = k(n)$? Here $k = k(n)$ has to be an increasing function of n .*

Remark 6.5. A knot is called n -adjacent to the unknot if it is n -trivial and each set of crossings in the definition of n -triviality has cardinality one. The

first named author of this paper and N. Askitas have proved Conjecture 6.3 for n -adjacent knots. ([A-K]).

REFERENCES

- [A-K] E. Kalfagianni and N. Askitas: *On n -adjacent knots*, preprint in preparation.
- [Bi] J. S. Birman: *New points of view in knot theory*, Bull. of AMS, **28**(1993), 253–287.
- [BN1] D. Bar-Natan: *On the Vassiliev knot invariants*, Topology, **34**(1995), 423–472.
- [BN2] D. Bar-Natan: *Vassiliev homotopy string link invariants*, J. of Knot Theory and Ramifications, **4**(1995), 13–32.
- [BN-G] D. Bar-Natan and S. Garoufalidis: *On the Melvin-Morton-Rozansky conjecture*, Invent. Math., **125**(1996), 103–133.
- [Co] T. D. Cochran: *Derivatives of Links: Milnor’s Concordance Invariants and Massey’s Products*, Memoirs AMS, **84** (1990).
- [C] J. Conant: Private communication.
- [C1] J. Conant: *A knot bounding a grope of class n is $n/2$ -trivial*, UCSD preprint 1999.
- [Gu] M. N. Gusarov: *On n -equivalence of knots and invariants of finite degree*, in *Topology of Manifolds and Varieties*, 173–192, Adv. Soviet Math., vol. 18, AMS, 1994.
- [He] J. Hempel: *3-manifolds*, Annals of Mathematics Studies, vol. 86, Princeton University Press, 1976.
- [H-M] N. Habegger and G. Masbaum: *The Kontsevich integral and Milnor’s Invariants*, preprint 1997.
- [I-O] K. Igusa and K. Orr: *Links, pictures and the homology of nilpotent groups*, preprint, 1998.
- [KSM] A. Karras, W. Magnus and D. Solitar: *Combinatorial Group Theory*, Pure and Appl. Math., vol. XIII, Interscience, NY, 1966.
- [K-L] E. Kalfagianni and X.-S. Lin: *Milnor and finite type invariants of plat-closures*, Math. Research Letters, to appear .
- [L1] X.-S. Lin: *Power series expansions and invariants of links*, in *Geometric Topology*, 184–202, AMS/IP Studies in Advanced Mathematics, vol. 2, 1996.

- [L2] X.-S. Lin: *Null k -cobordant links in S^3* , Comm. Math. Helv., **66**(1991), 333-339.
- [Mi1] J. Milnor: *Link groups*, Annals of Math., **2** (1954), 145-154.
- [Mi2] J. Milnor: *Isotopy of links*, in *Algebraic Geometry and Topology*, 280–386, Princeton University Press, 1957.
- [M-M] P. Melvin and H. Morton: *The colored Jones function*, Comm. Math. Phys., **169**(1995), 501-520.
- [N-S] K.-Y. Ng and T. Stanford: *On Gussarov's groups of knots*, Math. Proc. Camb. Phil. Soc., to appear.
- [O] K. Orr: *Homotopy invariants of links*, Invent. Math., **95**(1989), 379-394.
- [Ro] D. Rolfsen: *Knots and Links*, Publish or Perish, 1976.
- [Ru] L. Rudolph: *A congruence between link polynomials*, Math. Proc. Camb. Phil. Soc., **107** (1990), 319-327.
- [Ya] M. Yamamoto: *Knots and spatial embeddings of the complete graph on four vertices*, Top. and Appl., **3**(1990), 291-298.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI 48824

E-mail address: *kalfagia@math.msu.edu*

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, CA 92521

E-mail address: *xl@math.ucr.edu*